

Mirror symmetry, Tyurin degenerations and fibrations on Calabi-Yau manifolds

Charles F. Doran, Andrew Harder, and Alan Thompson

ABSTRACT. We investigate a potential relationship between mirror symmetry for Calabi-Yau manifolds and the mirror duality between quasi-Fano varieties and Landau-Ginzburg models. More precisely, we show that if a Calabi-Yau admits a so-called Tyurin degeneration to a union of two Fano varieties, then one should be able to construct a mirror to that Calabi-Yau by gluing together the Landau-Ginzburg models of those two Fano varieties. We provide evidence for this correspondence in a number of different settings, including Batyrev-Borisov mirror symmetry for K3 surfaces and Calabi-Yau threefolds, Dolgachev-Nikulin mirror symmetry for K3 surfaces, and an explicit family of threefolds that are not realized as complete intersections in toric varieties.

1. Introduction

The aim of this paper is to investigate the relationship between mirror symmetry for Calabi-Yau manifolds and for Fano varieties of the same dimension.

Classically, mirror symmetry is a network of conjectures relating the properties of two mirror dual Calabi-Yau manifolds. For us, unless otherwise stated, a *Calabi-Yau manifold* will always be a smooth compact Kähler manifold V with trivial canonical bundle $\omega_V \cong \mathcal{O}_V$ and vanishing cohomology groups $H^i(V, \mathcal{O}_V)$ for all $0 < i < \dim(V)$.

A similar duality has been proposed for Fano varieties. In physics, Eguchi, Hori and Xiong [EHX97] postulated that a d -dimensional manifold X with $c_1(X) > 0$ should be mirror to a *Landau-Ginzburg model* (Y, w) , where Y is a d -dimensional Kähler manifold and w is a function $w: Y \rightarrow \mathbb{C}$. This correspondence was then incorporated into the framework of homological mirror symmetry, as a correspondence between the directed Fukaya category [Sei01b] (resp. the bounded derived

2010 *Mathematics Subject Classification.* Primary 14J33; Secondary 14J32, 14J30, 14J28, 14D06.

Key words and phrases. Mirror symmetry, Calabi-Yau threefold, K3 surface, fibration, degeneration.

The first author would like to acknowledge support from the Natural Sciences and Engineering Research Council of Canada (NSERC) and the Visiting Campobassi Professorship of the University of Maryland.

The second author was supported by an NSERC Post-Graduate Scholarship.

The third author was supported by a Fields-Ontario Postdoctoral Fellowship with funding provided by NSERC and the Ontario Ministry of Training, Colleges and Universities.

category of singularities [Orl09]) associated to (Y, \mathbf{w}) and the bounded derived category of coherent sheaves on X (resp. the Fukaya category of X). More recently, Katzarkov, Kontsevich and Pantev [KKP14] conjectured that if X is a Fano variety, then the Landau-Ginzburg model (Y, \mathbf{w}) of X is in fact a quasi-projective variety that satisfies certain specific conditions and, moreover, that there is a mirror relationship between the Hodge numbers of X and certain Hodge-theoretic invariants of (Y, \mathbf{w}) .

It is expected that this notion of mirror symmetry for Fano varieties is related to classical mirror symmetry for Calabi-Yau manifolds of one dimension lower, as follows. If X is a d -dimensional Fano variety with mirror Landau-Ginzburg model (Y, \mathbf{w}) , then a general fibre of \mathbf{w} is expected to be a $(d-1)$ -dimensional Calabi-Yau variety that is mirror, in the classical sense, to a generic anticanonical hypersurface in X .

This raises a natural question: is mirror symmetry for d -dimensional Fano varieties related to classical mirror symmetry for Calabi-Yau manifolds of the same dimension d ? In this paper we outline a correspondence that provides a potential answer to this question.

This correspondence may be described as follows. Let V be a d -dimensional Calabi-Yau manifold and suppose that V admits a degeneration to a union $X_1 \cup_Z X_2$ of two quasi-Fano varieties glued along an anticanonical hypersurface Z (such degenerations are called *Tyurin degenerations*). Then we claim that the mirror W of V admits a fibration by $(d-1)$ -dimensional Calabi-Yau manifolds, with general fibre S that is mirror to Z . Moreover, W can be constructed topologically by gluing together the Landau-Ginzburg models (Y_1, \mathbf{w}_1) and (Y_2, \mathbf{w}_2) of X_1 and X_2 , in a sense to be made precise in Section 2.2.

The first person to observe traces of such a correspondence was probably Dolgachev [Dol96], who noticed that Dolgachev-Nikulin mirror symmetry for K3 surfaces matches Type II degenerations (of which Tyurin degenerations are a special case) with elliptic fibrations on the mirror. After this, the first mention of a higher dimensional version appears to be due to Tyurin, who gave a brief hint of its existence at the very end of [Tyu04].

More recently, a variant of the construction presented here was worked out in detail by Auroux [Aur08], in the special case where V is a double cover of a Fano variety X ramified over a smooth member of $| -2K_X |$; this V admits a Tyurin degeneration to the union of X with itself.

The structure of this paper is as follows. In Section 2 we describe our construction. We begin with a d -dimensional Calabi-Yau manifold V which admits a Tyurin degeneration to a union $X_1 \cup_Z X_2$ of quasi-Fano varieties glued along an anticanonical hypersurface Z . Then we show that the Landau-Ginzburg models (Y_1, \mathbf{w}_1) and (Y_2, \mathbf{w}_2) of X_1 and X_2 may be glued together to form a new variety W , which is fibred by Calabi-Yau $(d-1)$ -folds topologically mirror to Z , so that the Euler numbers of V and W satisfy the mirror relationship $\chi(V) = (-1)^d \chi(W)$. This suggests that V and W should be thought of as mirror dual. In the threefold case we provide even more evidence for this conjecture: if we make the assumptions that W is Calabi-Yau and that the K3 surface Z is Dolgachev-Nikulin mirror to a general fibre of the fibration on W , then we can show that V and W are in fact topologically mirror.

In the remaining sections of the paper we discuss this correspondence in several special cases. In Section 3 we discuss the case of Batyrev-Borisov mirror symmetry for surfaces and threefolds. Indeed, suppose that V is a K3 surface or Calabi-Yau threefold constructed as an anticanonical hypersurface in a Gorenstein toric Fano 3- or 4-fold, determined by a reflexive polytope Δ . We show that a nef partition Δ_1, Δ_2 of Δ determines both a Tyurin degeneration $X_1 \cup_Z X_2$ of V and a fibration $\pi: W \rightarrow \mathbb{P}^1$ on a birational model W of its Batyrev mirror, so that the general fibre of π is Batyrev-Borisov mirror dual to the intersection $Z = X_1 \cap X_2$.

Specializing to the threefold case, we further show that the singular fibres of the K3 surface fibration $\pi: W \rightarrow \mathbb{P}^1$ contain numerical information about X_1 and X_2 , and describe a relationship between W and the Landau-Ginzburg models of X_1 and X_2 . Unfortunately a corresponding result in the K3 surface case is difficult to prove for combinatorial reasons, but we conjecture the form that such a result should take.

Section 4 is concerned with Dolgachev-Nikulin mirror symmetry for K3 surfaces. We revisit Dolgachev's [Dol96] mirror correspondence between Type II degenerations (of which Tyurin degenerations are a special case) and elliptic fibrations, which may be thought of as a generalization of the correspondence described in Section 2. Consideration of several explicit examples suggests a way to enhance our conjectures to cope with more general Type II degenerations, which may contain more than two components.

In Section 5 we discuss how this theory fits with classical mirror symmetry for threefolds. We begin by showing that, if V is a Calabi-Yau threefold that undergoes a Tyurin degeneration (satisfying certain technical conditions), then mirror symmetry predicts the existence of a K3 fibration on the mirror threefold W , with properties consistent with those expected from the theory in Section 2. Following this, we specialize our discussion to the case of threefolds fibred by quartic mirror K3 surfaces, as studied in [DHNT16]. In this setting we explicitly construct candidate mirror threefolds, along with Tyurin degenerations of them, and show that they have the properties predicted by Section 2. In particular, this provides an important illustration of our theory using threefolds that are not complete intersections in toric varieties, thereby giving evidence that the ideas of Section 2 apply beyond the toric setting of Section 3.

Finally, Section 6 discusses the limitations of our construction. Indeed, it appears that difficulties arise for Tyurin degenerations of V which occur along loci in moduli that are disjoint from points of maximally unipotent monodromy. In this case, we seem to have no guarantee of the existence of a mirror fibration on W ; an example where this occurs is given in Example 6.1. Instead we present evidence that, if W is replaced by its bounded derived category of coherent sheaves $\mathbf{D}^b(W)$, it should be possible to find a non-commutative fibration of $\mathbf{D}^b(W)$ by Calabi-Yau categories, which might be thought of as homologically mirror to the Tyurin degeneration of V .

2. Setup and preliminary results

Our aim is to provide evidence for a mirror correspondence between a certain type of degeneration of Calabi-Yau manifolds, called a Tyurin degeneration, and Calabi-Yau manifolds constructed by gluing Landau-Ginzburg models. We begin by defining these objects.

2.1. Smoothing Tyurin degenerations. A smooth variety X is called a *quasi-Fano* variety if its anticanonical linear system contains a smooth Calabi-Yau member and $H^i(X, \mathcal{O}_X) = 0$ for all $i > 0$. Given this, a *Tyurin degeneration* is a degeneration $\mathcal{V} \rightarrow \Delta$ of Calabi-Yau manifolds over the unit disc $\Delta \subset \mathbb{C}$, such that the total space \mathcal{V} is smooth and the central fibre is a union of two quasi-Fano varieties that meet normally along a smooth variety Z , with $Z \in |-K_{X_i}|$ for each $i \in \{1, 2\}$. Degenerations of this type have been studied by Lee [Lee06], who coined the name.

This construction can be reversed, and a family of Calabi-Yau manifolds built up from a pair of quasi-Fano varieties X_1 and X_2 as follows. Let Z be a smooth variety which is a member of both $|-K_{X_1}|$ and $|-K_{X_2}|$, and suppose that there are ample classes $D_1 \in \text{Pic}(X_1)$ and $D_2 \in \text{Pic}(X_2)$ which both restrict to the same ample class $D \in \text{Pic}(Z)$ (this last condition is needed to ensure that [KN94, Theorem 4.2], which gives the existence of a smoothing, can be applied in our setting). Let $X_1 \cup_Z X_2$ denote the variety which is a normal crossings union of X_1 and X_2 meeting along Z .

With this setup, we say that $X_1 \cup_Z X_2$ is *smoothable* to a Calabi-Yau manifold V if there exists a complex manifold \mathcal{V} equipped with a map $\psi: \mathcal{V} \rightarrow \Delta$ so that the fibre $\psi^{-1}(0) = X_1 \cup_Z X_2$, the fibre $\psi^{-1}(t)$ is a smooth Calabi-Yau manifold for any $t \in \Delta \setminus \{0\}$, and V is a general fibre of \mathcal{V} . It follows from a theorem of Kawamata and Namikawa [KN94, Theorem 4.2] that $X_1 \cup_Z X_2$ is smoothable to a Calabi-Yau manifold V if and only if N_{Z/X_1} and N_{Z/X_2} are inverses of one another and, moreover, that the resulting manifold V is unique up to deformation.

2.2. Gluing Landau-Ginzburg models. Let us first define what we mean by Landau-Ginzburg (LG) model in this paper. In [Har16], a notion of a LG model is defined which conjecturally encapsulates the LG models of Fano varieties, and even goes further to describe the LG models of many quasi-Fano varieties. For general quasi-Fano varieties, however, we do not believe that this definition is sufficient; in particular, it seems that for general quasi-Fanos we must drop any expectation that our LG model be algebraic.

For this reason, in this paper we adopt a much more general definition.

DEFINITION 2.1. A *Landau-Ginzburg (LG) model* of a quasi-Fano variety is a pair (Y, w) consisting of a Kähler manifold Y satisfying $h^1(Y) = 0$ and a proper map $w: Y \rightarrow \mathbb{C}$. The map w is called the *superpotential*.

Note that this definition leaves room for the image of w to be an open set in \mathbb{C} . If Y is quasi-projective then the Hodge numbers of such LG models (Y, w) are defined in [KKP14]; however, in the general case it is unclear how this should be done. Instead, following [KKP14], we propose that if (Y, w) is the LG model of a quasi-Fano variety X , then we should have

$$(2.1) \quad h^i(Y, w^{-1}(t)) = \sum_j h^{d-i+j, j}(X),$$

where $h^i(Y, w^{-1}(t))$ is the rank of the cohomology group of the pair $H^i(Y, w^{-1}(t))$ and t is a generic point in the image of w . We also expect that if (Y, w) is the LG model of X , then the smooth fibres of w should be mirror to generic anticanonical hypersurfaces in X .

With notation as in the previous section, it now seems pertinent to ask whether there is any relationship between the LG models of the quasi-Fano varieties X_1 and X_2 , and mirror symmetry for V . Indeed, it seems natural to expect that these LG models could be somehow glued together to give a mirror W for V , since we are, in a topological sense, gluing X_1 and X_2 together to form V (see [Tyu04] for details on this topological construction).

In more detail, we expect that if Y_i is the LG model of X_i , equipped with superpotential w_i , then the monodromy symplectomorphism on $w_i^{-1}(t)$ (for t a regular value of w_i) associated to a small loop around ∞ can be identified under mirror symmetry with the restriction of the Serre functor of the bounded derived category of coherent sheaves $\mathbf{D}^b(X_i)$ on X_i to the bounded derived category of coherent sheaves $\mathbf{D}^b(Z)$ on Z [Sei01a, KKP14]. This Serre functor is simply $(-)\otimes\omega_{X_i}[d]$ where $[d]$ denotes shift by $d = \dim X_i$. Thus, up to a choice of shift, we see that the action of monodromy on $w_i^{-1}(t)$ should be identified with the autoequivalence of $\mathbf{D}^b(Z)$ induced by taking the tensor product with $\omega_{X_i}|_Z = N_{Z/X_i}^{-1}$.

Now recall that if $X_1 \cup_Z X_2$ is smoothable to V , then we have $N_{Z/X_1} \otimes N_{Z/X_2} = \mathcal{O}_Z$, so the monodromy symplectomorphism ϕ_1 associated to a clockwise loop around infinity on $w_1^{-1}(t)$ should be same as the monodromy ϕ_2^{-1} associated to a *counter-clockwise* loop around infinity on $w_2^{-1}(t)$. It should be noted that, for this to make sense, we must assume that the fibres of w_1 and w_2 are topologically the same Calabi-Yau manifold, which we denote by S ; this assumption is stronger than the assumption that both are mirror to Z .

Now we glue these LG models as follows. For each $i \in \{1, 2\}$, choose r_i so that $|\lambda| \leq r_i$ for every λ in the critical locus of w_i . Then choose local trivializations of Y_i over $U_i = \{z \in \mathbb{C} : |z| > r_i\}$ and let $Q_i = w_i^{-1}(U_i)$. This local trivialization is topologically equivalent to expressing Q_i as a gluing of the ends of $B_i = S \times [-1, 1] \times (-1, 1)$ together via the map

$$\tilde{\phi}_i: p \times \{-1\} \times (z) \mapsto \phi_i(p) \times \{1\} \times (z),$$

where ϕ_i is the monodromy symplectomorphism, and we identify $S \times \{-1\} \times (-1, 1)$ with $S \times \{1\} \times (-1, 1)$.

Assuming that $\phi_1 = \phi_2^{-1}$ (which, we recall, conjecturally follows from smoothability of $X_1 \cup_Z X_2$), we can identify B_1 with B_2 by the map

$$\tau: p \times [x] \times (z) \mapsto p \times [-x] \times (-z).$$

Under this identification of B_1 and B_2 , it is clear that $\tau \cdot \tilde{\phi}_1 = \tilde{\phi}_2$. Thus the identification τ gives an isomorphism between Q_1 and Q_2 , allowing us to glue Y_1 to Y_2 along Q_1 and Q_2 to produce a C^∞ manifold W . This gluing respects the fibrations w_1 and w_2 , so W is equipped with a fibration π over the gluing of \mathbb{C} with \mathbb{C} described above. It is clear that the base of this fibration is just the 2-sphere S^2 . This procedure is illustrated in Figure 1.

EXAMPLE 2.2. As a sanity check, we can perform this construction with elliptic curves. Take a degeneration of an elliptic curve to a union of two copies of \mathbb{P}^1 meeting in two points (Kodaira type I_2). The LG model of \mathbb{P}^1 is the map from \mathbb{C}^\times , which is topologically a twice-punctured rational curve, to \mathbb{C} given by

$$w: x \mapsto x + \frac{1}{x}.$$




FIGURE 1. Gluing Y_1 and Y_2 to give W

This map w is a double covering of \mathbb{A}^1 ramified at two points. One can check that monodromy of this fibration around the point at infinity is trivial. Let Y_1 and Y_2 be copies of this LG model of \mathbb{P}^1 . Then we may glue Y_1 and Y_2 as described above. The resulting topological space is a double cover of S^2 which is ramified at four points. This is simply the 2-dimensional torus, which is topologically mirror to the original elliptic curve

THEOREM 2.3. *Let X_1 and X_2 be d -dimensional quasi-Fano varieties which contain the same anticanonical Calabi-Yau hypersurface Z , such that $K_{X_1}|_Z + K_{X_2}|_Z = 0$. Let (Y_1, w_1) and (Y_2, w_2) be Landau-Ginzburg models of X_1 and X_2 , and suppose that the fibres of w_1 and w_2 are topologically the same Calabi-Yau manifold, which is topologically mirror to Z . Finally, let V be a Calabi-Yau variety obtained from $X_1 \cup_Z X_2$ by smoothing and let W be the variety obtained by gluing Y_1 to Y_2 as above. Then*

$$\chi(V) = (-1)^d \chi(W),$$

where χ denotes the Euler number.

PROOF. Start by recalling the long exact sequence of the pair $(Y_i, w_i^{-1}(t))$, for t a regular value of w_i ,

$$\dots \rightarrow H^n(Y_i, \mathbb{C}) \rightarrow H^n(w_i^{-1}(t), \mathbb{C}) \rightarrow H^{n+1}(Y_i, w_i^{-1}(t); \mathbb{C}) \rightarrow H^{n+1}(Y_i, \mathbb{C}) \rightarrow \dots$$

Since Euler numbers are additive in long exact sequences, we have that $\chi(Y_i) = \chi(Y_i, w_i^{-1}(t)) + \chi(w_i^{-1}(t))$. By Equation (2.1), we see that $\chi(Y_i, w_i^{-1}(t))$ is equal to $(-1)^d \chi(X_i)$, where d is the dimension of Y_i . Thus $\chi(Y_i) = (-1)^d \chi(X_i) + \chi(w_i^{-1}(t))$. Moreover, since $w_i^{-1}(t)$ is topologically mirror to Z by assumption, we have that $\chi(Z) = (-1)^{d-1} \chi(w_i^{-1}(t))$, which gives $\chi(Y_i) = (-1)^d (\chi(X_i) - \chi(Z))$.

On the other hand, the Mayer-Vietoris exact sequence

$$\dots \rightarrow H^n(W, \mathbb{C}) \rightarrow H^n(Y_1, \mathbb{C}) \oplus H^n(Y_2, \mathbb{C}) \rightarrow H^n(Y_1 \cap Y_2, \mathbb{C}) \rightarrow \dots$$

gives $\chi(W) = \chi(Y_1) + \chi(Y_2) - \chi(Y_1 \cap Y_2)$. Since $Y_1 \cap Y_2$ is a fibration over an annulus, we can compute its cohomology using the Wang sequence [PS08, Theorem 11.33]

$$\cdots \rightarrow H^n(Y_1 \cap Y_2, \mathbb{C}) \rightarrow H^n(w^{-1}(t), \mathbb{C}) \xrightarrow{T_n - \text{Id}} H^n(w^{-1}(t), \mathbb{C}) \rightarrow \cdots,$$

where T_n is the action of monodromy on $H^n(w^{-1}(t), \mathbb{C})$ associated to a small loop around our annulus, to obtain $\chi(Y_1 \cap Y_2) = 0$. Putting everything together, we obtain

$$\chi(W) = (-1)^d(\chi(X_1) + \chi(X_2) - 2\chi(Z)).$$

Finally, since $X_1 \cup_Z X_2$ is smoothable to V , we can compute the Euler characteristic of V by applying [Lee06, Proposition IV.6], which states that

$$\chi(V) = \chi(X_1) + \chi(X_2) - 2\chi(Z).$$

We therefore have that $\chi(W) = (-1)^d\chi(V)$, as claimed. \square

This is precisely the relationship between the Euler characteristics of mirror dual Calabi-Yau varieties. In the next subsection, we will provide more evidence for the hypothesis that W is the mirror dual of the original Calabi-Yau variety V , in the special case where V is a Calabi-Yau threefold.

REMARK 2.4. Note that the requirement that there exist two ample divisors D_1 and D_2 , on X_1 and X_2 respectively, which restrict to the same divisor on Z was not used at all in the proof of Theorem 2.3. Moreover, despite the fact that the proof of [KN94, Theorem 4.2] uses this assumption in a material way (in order to prove the pro-representability of the log deformation functor), the topological construction of the gluing of X_1 and X_2 can be performed without it.

For instance, let us take a generic K3 surface Z with Picard lattice of rank 2 isomorphic to the lattice with Gram matrix

$$\begin{pmatrix} 4 & 6 \\ 6 & 6 \end{pmatrix}.$$

Such a K3 surface embeds into both \mathbb{P}^3 and the intersection of a quadric Q and a cubic C in \mathbb{P}^5 . Let us blow up \mathbb{P}^3 in $Z \cap Z'$ for some generic K3 surface Z' in \mathbb{P}^3 , calling the result X_1 , and blow up $Q \cap C$ in the intersection of Z and a generic hyperplane section in \mathbb{P}^5 , calling the result X_2 . Then the normal crossings variety $X_1 \cup_Z X_2$ is not Kähler, so we cannot find D_1 and D_2 as above. However, both V and W can be constructed, as C^∞ manifolds, from $X_1 \cup_Z X_2$ by the method we have described. We wonder whether V and W represent a mirror pair of non-Kähler Calabi-Yau manifolds.

2.3. The threefold case. With notation as before, Lee [Lee10] has computed the Hodge numbers of V in the case where X_1 and X_2 are smooth threefolds. Let us define $\rho_i: H^2(X_i, \mathbb{Q}) \rightarrow H^2(Z, \mathbb{Q})$ for $i = 1, 2$ to be the restriction and define $k = \text{rank}(\text{im}(\rho_1) + \text{im}(\rho_2))$.

THEOREM 2.5. [Lee10, Corollary 8.2] *Let V be a Calabi-Yau threefold constructed as a smoothing of $X_1 \cup_Z X_2$, as above. Then*

$$\begin{aligned} h^{1,1}(V) &= h^2(X_1) + h^2(X_2) - k - 1, \\ h^{2,1}(V) &= 21 + h^{2,1}(X_1) + h^{2,1}(X_2) - k. \end{aligned}$$

On the other side of the picture, we have a corresponding result for W .

PROPOSITION 2.6. *Let W be as above and let S be a general fibre of the map π . Assume that $\dim W = \dim S + 1 = 3$. Then*

$$h^2(W) = 1 + h^2(Y_1, S) + h^2(Y_2, S) + \ell,$$

where ℓ is the rank of the subgroup of $H^2(S, \mathbb{C})$ spanned by the intersection of the images of $H^2(Y_1, \mathbb{C})$ and $H^2(Y_2, \mathbb{C})$ under the natural restriction maps.

PROOF. Let U be the annulus along which B_1 and B_2 are glued, and let $Q = \pi^{-1}(U)$ be its preimage in W . We begin by computing the rank of $H^2(W, \mathbb{C})$ using the Mayer-Vietoris sequence

$$\cdots \rightarrow H^1(Q, \mathbb{C}) \rightarrow H^2(W, \mathbb{C}) \rightarrow H^2(Y_1, \mathbb{C}) \oplus H^2(Y_2, \mathbb{C}) \xrightarrow{r_1^Q - r_2^Q} H^2(Q, \mathbb{C}) \rightarrow \cdots,$$

where r_i^Q are the natural restriction maps from $H^2(Y_i, \mathbb{C})$ to $H^2(Q, \mathbb{C})$. From the Wang sequence, we obtain $H^1(Q, \mathbb{C}) = \mathbb{C}$. So, using the assumption that $H^1(Y_1, \mathbb{C}) = H^1(Y_2, \mathbb{C}) = 0$, we see that $H^2(W, \mathbb{C})$ is isomorphic to the direct product of \mathbb{C} and the kernel of the restriction map $r_1^Q - r_2^Q$. We note that this map fits into a commutative triangle

$$\begin{array}{ccc} H^2(Y_1, \mathbb{C}) \oplus H^2(Y_2, \mathbb{C}) & \xrightarrow{r_1^Q - r_2^Q} & H^2(Q, \mathbb{C}) \\ & \searrow r_1^S - r_2^S & \downarrow r_Q^S \\ & & H^2(S, \mathbb{C}). \end{array}$$

Now, since S is a K3 surface, we have $h^1(S) = 0$, and it follows from the Wang sequence that the map r_Q^S is injective. Thus the kernel of $r_1^S - r_2^S$ is the same as the kernel of $r_1^Q - r_2^Q$. Elementary linear algebra gives that the rank of this kernel is $h^2(Y_1) + h^2(Y_2) - \text{rank}(\text{im}(r_1^S) + \text{im}(r_2^S))$. So we obtain

$$h^2(W) = 1 + h^2(Y_1) + h^2(Y_2) - \text{rank}(\text{im}(r_1^S) + \text{im}(r_2^S)).$$

Now, for $i = 1, 2$ we have exact sequences

$$0 \longrightarrow H^2(Y_i, S; \mathbb{C}) \longrightarrow H^2(Y_i, \mathbb{C}) \xrightarrow{r_i^S} H^2(S, \mathbb{C}) \longrightarrow \cdots$$

Which give

$$h^2(Y_i) = h^2(Y_i, S) + \text{rank}(\text{im}(r_i^S)).$$

Putting together with the previous expression, the proposition follows. □

Therefore, if W admits a complex structure for which it is Calabi-Yau, then we compute

$$\begin{aligned} \chi(W) &= 2h^{1,1}(W) - 2h^{2,1}(W) \\ &= 2(1 + h^2(Y_1, S) + h^2(Y_2, S) + \ell) - 2h^{2,1}(W). \end{aligned}$$

Equation (2.1) then gives that $h^2(Y_i, S) = h^{2,1}(X_i)$, so

$$\chi(W) = 2(h^{2,1}(X_1) + h^{2,1}(X_2) - h^{2,1}(W) + \ell + 1).$$

Furthermore, from Theorems 2.3 and 2.5, we also know that

$$\begin{aligned}\chi(W) &= -\chi(V) \\ &= -2h^{1,1}(V) + 2h^{2,1}(V) \\ &= -2(h^2(X_1) + h^2(X_2) - k - 1) + 2(h^{2,1}(X_1) + h^{2,1}(X_2) + 21 - k) \\ &= 2(h^{2,1}(X_1) + h^{2,1}(X_2) - h^2(X_1) - h^2(X_2) + 22)\end{aligned}$$

Putting this together, we have that $h^{2,1}(W) = \ell - 21 + h^2(X_1) + h^2(X_2)$. So in order for W and V to be topologically mirror to one another, we must have

$$\ell - 21 + h^2(X_1) + h^2(X_2) = h^2(X_1) + h^2(X_2) - k - 1,$$

which is equivalent to $\ell + k = 20$. This is true if S and Z are mirror dual in the sense of Dolgachev-Nikulin, given the lattice polarization on Z (resp. S) coming from the sum of the images of the restriction maps $H^2(X_i, \mathbb{Z}) \rightarrow H^2(Z, \mathbb{Z})$ (resp. the intersection of the images of the restriction maps $H^2(Y_i, \mathbb{Z}) \rightarrow H^2(S, \mathbb{Z})$). Thus mirror symmetry for V and W is consistent with mirror symmetry for S and Z .

REMARK 2.7. In the case where W is Calabi-Yau and S and Z are Dolgachev-Nikulin mirror dual, the expressions

$$\begin{aligned}h^{2,1}(V) &= 21 + h^{2,1}(X_1) + h^{2,1}(X_2) - k \\ h^{1,1}(W) &= 1 + h^2(Y_1, S) + h^2(Y_2, S) + \ell\end{aligned}$$

could be thought of as mirror dual decompositions of the corresponding Hodge numbers, in the following sense.

The Hodge number $h^{2,1}(X_i)$ may be interpreted as the fibre dimension of the natural map from the moduli space of pairs (X_i, Z) to the moduli space of appropriately polarized K3 surfaces Z (which has dimension $20 - k$). Thus the degenerate fibre $X_1 \cup_Z X_2$ should have $h^{2,1}(X_1) + h^{2,1}(X_2) + 20 - k = h^{2,1}(V) - 1$ deformations, and such Tyurin degenerations should appear in codimension 1 in the moduli space of V .

We thus obtain a decomposition of $h^{2,1}(V)$ into contributions $h^{2,1}(X_i)$ coming from deformations of each X_i , a contribution $(20 - k)$ from deformations of the gluing locus Z , and 1 for the codimension in the moduli space.

On the mirror side a similar statement holds: $h^{1,1}(W)$ can be decomposed into contributions $h^2(Y_i, S)$ coming from the LG-models (Y_i, w_i) (these will be interpreted later as counts of components in singular fibres), a contribution ℓ from divisors on the generic fibre S , and 1 for the class of a general fibre (compare [DHNT16, Lemma 3.2]). The picture is completed by noting that $h^2(Y_i, S) = h^{2,1}(X_i)$ and $\ell = 20 - k$.

3. Batyrev-Borisov mirror symmetry

In this section we will prove a number of results that illustrate the situation considered in the previous section in the special case of Batyrev-Borisov mirror symmetry. For background on the definitions and concepts used in this section, we refer the reader to [CK99, CLS11]. However, since our conventions differ very slightly from those used in the references above, before we proceed we will briefly outline the notation to be used in the remainder of this section.

Let M be a free \mathbb{Z} -module of rank d , let Δ be a reflexive polytope in $M \otimes \mathbb{R} = M_{\mathbb{R}}$, and denote the boundary of Δ by $\partial\Delta$. Let $N = \text{Hom}(M, \mathbb{Z})$ be the dual lattice to M and denote by $\langle \cdot, \cdot \rangle$ the natural bilinear pairing from $N \times M$ to \mathbb{Z} . Let

$$\Delta^\circ = \{u \in N_{\mathbb{R}} : \langle u, v \rangle \geq -1 \text{ for all } v \in \Delta\}$$

denote the polar polytope to Δ .

Let \mathbb{P}_Δ be the d -dimensional toric variety associated to the polytope Δ . The toric variety \mathbb{P}_Δ is Fano and has at worst Gorenstein singularities. Following [Bat94, Theorem 2.2.24], one can find a toric variety X_Δ which is a toric partial resolution of singularities of \mathbb{P}_Δ and which has at worst Gorenstein terminal singularities. Such X_Δ is referred to as a *maximal projective crepant partial (mpcp) resolution of singularities* of \mathbb{P}_Δ . In the future, we shall fix one such X_Δ for any given \mathbb{P}_Δ . The variety X_Δ can be presented as a quotient of some Zariski open subset $U \subseteq \mathbb{C}^{|\partial\Delta \cap M|}$ by the torus $(\mathbb{C}^\times)^{|\partial\Delta| - d}$. There is thus a homogeneous coordinate ring $\mathbb{C}[\{z_\rho\}_{\rho \in \partial\Delta \cap M}]$ on X_Δ .

The vanishing of each coordinate z_ρ determines a divisor on X_Δ , invariant under the natural action of the torus $(\mathbb{C}^\times)^d$, which we call D_ρ . The anticanonical divisor $-K_{X_\Delta}$ of X_Δ is linearly equivalent to $\sum_{\rho \in \partial\Delta \cap M} D_\rho$, and the cycle class group $A_1(X_\Delta)$ is generated by the divisors D_ρ . A divisor $\sum_{\rho \in \partial\Delta \cap M} b_\rho D_\rho$ for $b_\rho \in \mathbb{Z}$ is Cartier if and only if there is a piecewise linear function φ on $M_{\mathbb{R}}$, which takes integral values on M and which is linear on the cones of the fan defining X_Δ , so that $\varphi(\rho) = b_\rho$ for all ρ .

A *nef partition* of Δ is a partition of $\partial\Delta \cap M$ into sets E_1, \dots, E_k , so that for each $i = 1, \dots, k$, the divisor $\sum_{\rho \in E_i} D_\rho$ is nef and Cartier. Let us denote the line bundle thus associated to E_i by \mathcal{L}_i . We will let $\Delta_i = \text{Conv}(E_i \cup \{0_M\})$; in a mild abuse of terminology, we also refer to $\Delta_1, \dots, \Delta_k$ as a nef partition of Δ .

Batyrev’s [Bat94] toric version of mirror symmetry claims that the generic anticanonical hypersurfaces in X_Δ and X_{Δ° are mirror dual. Moreover, if we have a nef partition of Δ , then the complete intersection V of generic sections of $\mathcal{L}_1, \dots, \mathcal{L}_k$ is again Calabi-Yau. Borisov [Bor93] and Batyrev-Borisov [BB96] propose that there is a similar combinatorial construction of the mirror of V . In this case, we define

$$\nabla_i = \left\{ u \in N_{\mathbb{R}} : \begin{array}{l} \langle u, v \rangle \geq 0 \text{ for all } v \in E_j, j \neq i \\ \langle u, v \rangle \geq -1 \text{ for all } v \in E_i \end{array} \right\}$$

and let $\nabla = \text{Conv}(\nabla_1 \cup \dots \cup \nabla_k)$. This is a reflexive polytope and $\nabla_1, \dots, \nabla_k$ is a nef partition of ∇ . The complete intersection W in X_∇ cut out by generic sections of the line bundles associated to $\nabla_1, \dots, \nabla_k$ is a Calabi-Yau variety, which is expected to be mirror dual to V .

Finally, a *refinement* of a nef partition E_1, \dots, E_k is defined to be another nef partition F_1, \dots, F_{k+1} so that $F_i = E_i$ for $1 \leq i \leq k - 1$ and $E_k = F_k \cup F_{k+1}$.

Now, let X_Δ be a d -dimensional toric variety as above. Suppose that V is a Calabi-Yau complete intersection of nef divisors in X_Δ , determined by a nef partition E_1, \dots, E_k . Our aim is to show that, if F_1, \dots, F_{k+1} is a refinement of E_1, \dots, E_k , then this combinatorial data determines

- a Tyurin degeneration of V , and
- a pencil of quasi-smooth varieties birational to Calabi-Yau $(d - k - 1)$ -folds inside of the Batyrev-Borisov mirror W .

In the case where V is a threefold, we show that this pencil induces a K3 surface fibration on some birational model of W and that the singular fibres of this fibration carry information about the Tyurin degeneration of V . We will then compare this with the LG model picture in the previous section.

3.1. Tyurin degenerations. More precisely, let \mathcal{L}_i be the line bundles on X_Δ associated to the E_i . The refinement F_1, \dots, F_{k+1} gives rise to a pair of nef line bundles \mathcal{L}'_k and \mathcal{L}'_{k+1} so that $\mathcal{L}'_k \otimes \mathcal{L}'_{k+1} = \mathcal{L}_k$. Let $s_i \in H^0(X_\Delta, \mathcal{L}_i)$ be generic sections determining a quasi-smooth Calabi-Yau complete intersection V in X_Δ . If we let s'_k and s'_{k+1} be sections of \mathcal{L}'_k and \mathcal{L}'_{k+1} respectively, then $s'_k s'_{k+1}$ is a section of \mathcal{L}_k .

We can use this to construct a pencil of complete intersections as follows. First, let

$$V' = \bigcap_{i=1}^{k-1} \{s_i = 0\}$$

and assume that V' is connected and quasi-smooth; it is also clear that V' is quasi-Fano. Then take the pencil

$$\mathcal{Q}: \{ts_k - s'_k s'_{k+1} = 0\} \cap V'$$

in $\mathbb{A}^1 \times X_\Delta$, with t a parameter on \mathbb{A}^1 . If we assume that X_Δ is a smooth resolution of \mathbb{P}_Δ , then the only singularities of \mathcal{Q} in a neighbourhood of $0 \in \mathbb{A}^1$ are along $t = s_k = s'_k = s'_{k+1} = 0$, which we call Σ .

Note that since Σ is the intersection of a set of nef divisors in X_Δ , it has no base locus and its singularities are contained in the singular set of X_Δ . Furthermore, the intersection of Σ with any torus invariant subvariety of X_Δ is irreducible, thus Σ itself is either irreducible or a union of non-intersecting subvarieties of X_Δ . As a result, if X_Δ is smooth, then so is Σ , for general enough choices of sections.

We can thus resolve the singularities of \mathcal{Q} by blowing up $t = s'_k = 0$ inside of $\mathbb{A}^1 \times X_\Delta$ and taking the proper transform of \mathcal{Q} . The result is a Tyurin degeneration of V , so that the fibre over 0 of the degeneration is equal to the union of \hat{X}_1 and X_2 , where \hat{X}_1 is a quasi-Fano variety given by blowing up $X_1 := V' \cap \{s'_k = 0\}$ along $V' \cap \{s_k = s'_k = s'_{k+1} = 0\}$ and X_2 is a quasi-Fano variety given by $X_2 := V' \cap \{s'_{k+1} = 0\}$.

In the general situation, where X_Δ is not a smooth resolution of \mathbb{P}_Δ , we can still perform all of the steps above, but we will have singularities occurring at every step in general. The resulting degeneration will not be a Tyurin degeneration in the strict sense, but should still include data corresponding to the quasi-Fano varieties X_1, X_2 and the blown up locus Σ . We note here that a version of the smoothability result of Kawamata and Namikawa that works for mildly singular varieties has been explored in the thesis of Lee [Lee06]. The singular case may also be interpreted as equipping the union of X_1 and X_2 with a log structure (see e.g. [ACG13] and the references therein), which accounts for the subvariety Σ and determines the smoothing to V .

3.2. Pencils and fibrations on the mirror. Now we will look at how this nef partition is reflected in the mirror. For ease of notation we restrict ourselves to the case where V is a hypersurface; all of the results below generalize in the obvious way to refinements of k -partite nef partitions corresponding to codimension k complete intersections.

Since V is a hypersurface, the nef partition $E_1 = \Delta$ is trivial. Let Δ_1, Δ_2 denote the polytopes corresponding to the refinement F_1, F_2 of E_1 ; we thus have $\Delta = \text{Conv}(\Delta_1 \cup \Delta_2)$.

Now, the Batyrev dual of V is a Calabi-Yau variety W embedded as an anti-canonical hypersurface in X_{Δ° . By definition, W is cut out by an equation in the homogeneous coordinate ring of X_{Δ° , which may be written as

$$f := \sum_{\rho \in \Delta \cap M} a_\rho \prod_{\sigma \in \partial \Delta^\circ \cap N} z_\sigma^{\langle \sigma, \rho \rangle + 1} = 0,$$

where a_ρ are generically chosen complex coefficients. We will take a pencil \mathcal{P} of hypersurfaces in W , for $[s : t] \in \mathbb{P}^1$, defined by the intersection of W with hypersurfaces of the form

$$s \sum_{\rho \in \Delta_1 \cap M \setminus 0_M} a_\rho \prod_{\sigma \in \partial \Delta^\circ \cap N} z_\sigma^{\langle \sigma, \rho \rangle + 1} = t a_0 \prod_{\sigma \in \partial \Delta^\circ \cap N} z_\sigma.$$

Note that, away from $[s : t] = [0 : 1]$, this pencil may also be defined by the pair of equations

$$\begin{aligned} s \sum_{\rho \in \Delta_1 \cap M \setminus 0_M} a_\rho \prod_{\sigma \in \partial \Delta^\circ \cap N} z_\sigma^{\langle \sigma, \rho \rangle + 1} &= t a_0 \prod_{\sigma \in \partial \Delta^\circ \cap N} z_\sigma \\ s \sum_{\rho \in \Delta_2 \cap M \setminus 0_M} a_\rho \prod_{\sigma \in \partial \Delta^\circ \cap N} z_\sigma^{\langle \sigma, \rho \rangle + 1} &= (s - t) a_0 \prod_{\sigma \in \partial \Delta^\circ \cap N} z_\sigma. \end{aligned}$$

as the sum of these two equations is just sf , which vanishes precisely along W if $s \neq 0$.

We wish to show that the pencil \mathcal{P} induces a fibration on W . If we can show that a general member of this pencil is Calabi-Yau, then this will follow from:

PROPOSITION 3.1. *If $S \subseteq W$ is a smooth Calabi-Yau $(d - 1)$ -fold in a d -dimensional Calabi-Yau manifold, then the linear system $|S|$ is base-point free and hence there is a map $\pi : W \rightarrow \mathbb{P}^1$ with S as a fibre.*

PROOF. By adjunction, $\mathcal{O}_S(S) = N_{S/W} = \omega_S = \mathcal{O}_S$. Let $\iota : S \hookrightarrow W$ be the inclusion map, then we have a short exact sequence of sheaves

$$0 \longrightarrow \mathcal{O}_W \xrightarrow{s} \mathcal{O}_W(S) \longrightarrow \iota^* \mathcal{O}_S \longrightarrow 0,$$

where s is a section of $\mathcal{O}_W(S)$ whose vanishing locus is S . This short exact sequence gives rise to a long exact sequence in cohomology

$$0 \longrightarrow H^0(W, \mathcal{O}_W) \longrightarrow H^0(W, \mathcal{O}_W(S)) \longrightarrow H^0(W, \iota^* \mathcal{O}_S) \longrightarrow 0,$$

where the vanishing of $H^1(W, \mathcal{O}_W)$ follows from the Calabi-Yau property of W . From this sequence, we see that the restriction of a generic section of $\mathcal{O}_W(S)$ to S is a nonzero section of $\iota^* \mathcal{O}_S$, which is *a fortiori* non-vanishing. Thus $|S|$ is base-point free and determines a map $W \rightarrow \mathbb{P}^1$, since $h^0(\mathcal{O}_W(S)) = 2$ by the exact sequence above. □

To apply this proposition, we need to show that a general member S of the pencil \mathcal{P} is Calabi-Yau. When $\dim(W) = 2$ or 3 , this will follow from the next proposition, which is proved in [Har16].

PROPOSITION 3.2. [**Har16**] *If $\dim(W) = 2$, then a general member of the pencil \mathcal{P} is a smooth elliptic curve. If $\dim(W) = 3$, then a general member of the pencil \mathcal{P} is a smooth blow-up of a K3 surface S and, moreover, S is Batyrev-Borisov dual to the intersection of the quasi-Fano varieties X_1 and X_2 from Section 3.1.*

In the case where $\dim(W) = 2$, it therefore follows immediately from Proposition 3.1 that the pencil \mathcal{P} is an elliptic fibration on W . However, if W is a Calabi-Yau threefold and a general member of \mathcal{P} is a blown up K3 surface \hat{S} , then we will need to get rid of the (-1) -curves in S before we can apply Proposition 3.1. We will do this by performing a series of birational transformations.

LEMMA 3.3. *Let W be a Calabi-Yau threefold and let \hat{S} be a blown up K3 surface in W . If C is a (-1) -curve in \hat{S} , then $N_{C/W} \cong \mathcal{O}_{\mathbb{P}^1}(-1) \oplus \mathcal{O}_{\mathbb{P}^1}(-1)$.*

PROOF. We have a short exact sequence of sheaves on C ,

$$0 \longrightarrow \Theta_C \longrightarrow \Theta_W|_C \longrightarrow N_{C/W} \longrightarrow 0.$$

Since $c_1(\Theta_C) = 2$ and $c_1(\Theta_W|_C) = c_1(\Theta_W)|_C = 0$, it follows that $N_{C/W} \cong \mathcal{O}_{\mathbb{P}^1}(a) \oplus \mathcal{O}_{\mathbb{P}^1}(b)$ for some $a, b \in \mathbb{Z}$ with $a + b = -2$ (see, for example, [**Kat92**, Section 1]).

We may embed the normal bundle $N_{C/\hat{S}}$ into $N_{C/W}$ to get a short exact sequence of line bundles

$$0 \longrightarrow N_{C/\hat{S}} \longrightarrow N_{C/W} \longrightarrow \mathcal{L} \longrightarrow 0$$

for some line bundle \mathcal{L} . Since C is a (-1) -curve in \hat{S} , we know that $N_{C/\hat{S}} \cong \mathcal{O}_{\mathbb{P}^1}(-1)$. Furthermore, we have that $c_1(\mathcal{L}) = -1$ from the fact that $c_1(N_{C/W}) = -2$, thus $\mathcal{L} \cong \mathcal{O}_{\mathbb{P}^1}(-1)$. The long exact sequence in cohomology coming from the above short exact sequence proves that $H^0(C, N_{C/W}) = 0$ and therefore we must have that $N_{C/W} = \mathcal{O}_{\mathbb{P}^1}(-1) \oplus \mathcal{O}_{\mathbb{P}^1}(-1)$. \square

Therefore, any (-1) -curve C in \hat{S} may be blown up to produce a variety \widetilde{W} with exceptional divisor a copy of $\mathbb{P}^1 \times \mathbb{P}^1$. This copy of $\mathbb{P}^1 \times \mathbb{P}^1$ can be smoothly contracted along either ruling. Contracting along one ruling recovers W ; we denote the variety obtained by contracting along the other ruling by W^+ . The strict transform of \hat{S} in \widetilde{W} is just \hat{S} itself, but the contraction $\widetilde{W} \rightarrow W^+$ contracts the (-1) -curve C in \hat{S} . This is an example of a birational operation called a *flop*.

Repeating this for all (-1) curves in \hat{S} , we obtain a birational model of W which is a smooth Calabi-Yau threefold in which \hat{S} has been contracted to its minimal model, which is a K3 surface. Call the Calabi-Yau threefold resulting from this process \hat{W} . Applying Proposition 3.1 to \hat{W} , we see that we have proved:

THEOREM 3.4. *Let W be a Calabi-Yau threefold containing a smooth blown up K3 surface \hat{S} . By performing a sequence of flops on W , we may obtain a birational model \hat{W} of W which admits a fibration $\pi: \hat{W} \rightarrow \mathbb{P}^1$, so that the minimal model S of \hat{S} is a general fibre of π .*

Putting everything together, in the case of toric hypersurface Calabi-Yau threefolds we obtain:

COROLLARY 3.5. *If Δ is a 4-dimensional reflexive polytope that admits a nef partition Δ_1, Δ_2 and V is a general anticanonical hypersurface in X_Δ , then V admits a Tyurin degeneration and its Batyrev dual W has a birational model which admits a fibration by K3 surfaces. Moreover, the general fibre in this fibration on*

W is Batyrev-Borisov mirror dual to the complete intersection K3 surface in X_Δ determined by Δ_1, Δ_2 .

REMARK 3.6. There seems to be an inherent incompatibility between Batyrev (and Batyrev-Borisov) duality and K3 surface fibrations on Calabi-Yau threefolds, since one can show that even in very basic examples, there are exceptional curves in \hat{S} that cannot be avoided by simply changing the birational model of the toric ambient space X_{Δ° . A notable exception occurs when either Δ_1 or Δ_2 is 1-dimensional. In this case, one of the two component quasi-Fano varieties to which V degenerates is itself a toric variety. This is mirrored by the fact that X_{Δ° admits a morphism to \mathbb{P}^1 which induces the required K3 fibration on W . These seem to be a subset of the “toric fibrations” which have been studied extensively in the physics literature by a number of authors ([AKMS97, CCS13, GP13] to name a few).

REMARK 3.7. One can consider more general refinements of nef partitions, by taking a nef partition F_1, \dots, F_ℓ so that for each E_i , there is a subset I_i of $\{1, \dots, \ell\}$ so that $E_i = \cup_{j \in I_i} F_j$. These will give rise to generalized degenerations of the Calabi-Yau V determined by E_1, \dots, E_k to unions of quasi-Fano varieties, and families of Calabi-Yau varieties of codimension $\ell - k$ in W . The issue here, of course, is that it is hard to prove that this gives a fibration on a birational model of W . Despite this, these families of Calabi-Yau varieties surely have properties related to the LG models of the appropriate quasi-Fano varieties.

REMARK 3.8. Generalizing Corollary 3.5 to higher dimensions seems to be a challenge, since we have made use of both the minimal model program for surfaces and a characterization of flops in three dimensions. Of course, both of these objects have analogues in higher dimensions, but they are much more oblique and not likely to be useful in such a general situation. As it stands, the results above suffice to show that we obtain rational maps from W to \mathbb{P}^1 corresponding to every bipartite nef partition of Δ .

3.3. Singular fibres and a comparison with LG models. Now we will analyze the singular members of the pencil \mathcal{P} , which are birational to the fibres of \hat{W} . Using this, we can give a very accurate description of the singular fibres of \hat{W} , up to birational transformations. As we shall see, the resulting theory meshes nicely with the LG model picture described in Section 2. The results in this section can be extended quite generally, but for simplicity we will restrict ourselves to the situation of threefold hypersurfaces.

With notation as in the previous section, the nef partition Δ_1, Δ_2 of Δ determines a pair of polytopes $\nabla_1, \nabla_2 \subseteq \Delta^\circ$, which are a nef partition of $\nabla := \text{Conv}(\nabla_1 \cup \nabla_2) \subseteq \Delta^\circ$. Note that this inclusion may be strict; it therefore does not follow that we have a degeneration of W to quasi-Fano varieties dual to X_1 and X_2 .

Our first goal is to look at the member of the pencil \mathcal{P} over $[s : t] = [0 : 1]$ and understand its meaning in terms of the Tyurin degeneration described in Section 3.1.

PROPOSITION 3.9. *The member of the pencil \mathcal{P} corresponding to $[s : t] = [0 : 1]$ is equal to*

$$\bigcup_{\sigma \in (\Delta^\circ \setminus \nabla) \cap N} (D_\sigma \cap W).$$

In other words, the linear system defining the pencil \mathcal{P} is associated to the line bundle $\mathcal{O}_W(\sum_{\sigma \in (\Delta^\circ \setminus \nabla) \cap N} D_\sigma|_W)$.

PROOF. Recall that the pencil \mathcal{P} is defined as the intersection of W with hypersurfaces of the form

$$s \sum_{\rho \in \Delta_1 \cap M \setminus 0_M} a_\rho \prod_{\sigma \in \partial \Delta^\circ \cap N} z_\sigma^{\langle \sigma, \rho \rangle + 1} - ta_0 \prod_{\sigma \in \partial \Delta^\circ \cap N} z_\sigma = 0.$$

Thus, at least on an open set of W , we may write this family of hypersurfaces as a rational map from X_{Δ° to \mathbb{P}^1 , defined by $\phi: [z_\sigma] \mapsto [s : t]$ where

$$s = \prod_{\sigma \in (\Delta^\circ \setminus \nabla) \cap N} z_\sigma$$

$$t = \left(\sum_{\rho \in \Delta_1 \cap M \setminus 0_M} a_\rho \prod_{\sigma \in \partial \Delta^\circ \cap N} z_\sigma^{\langle \sigma, \rho \rangle + 1} \right) / \left(\prod_{\sigma \in \partial \nabla \cap N} z_\sigma \right)$$

We want to show that this map is defined on W away from the base locus of \mathcal{P} . Note that homogeneity away from $\sum_{\sigma \in \partial \nabla \cap N} D_\sigma$ is clear, since both terms are sections of $\mathcal{L} := \mathcal{O}_{X_{\Delta^\circ}}(\sum_{\sigma \in (\Delta^\circ \setminus \nabla) \cap N} D_\sigma)$. Now, if $\sigma \in \nabla_2$, then the numerator in the expression for t above has a factor of z_σ , since $\langle \sigma, \rho \rangle \geq 0$ for all $\rho \in \Delta_1$ and $\sigma \in \nabla_2$. Thus both terms are also sections of \mathcal{L} along D_σ for $\sigma \in \nabla_2$. Moreover, on the restriction of ϕ to W , we notice that the expression for t can also be written as

$$\left(\sum_{\rho \in \Delta_2 \cap M} a_\rho \prod_{\sigma \in \partial \Delta^\circ \cap N} z_\sigma^{\langle \sigma, \rho \rangle + 1} \right) / \left(\prod_{\sigma \in \partial \nabla \cap N} z_\sigma \right).$$

and thus, for the same reason as above, both terms are sections of \mathcal{L} along $D_\sigma \cap W$ for $\sigma \in \nabla_1$. Finally, [BB96, Proposition 6.3] implies that $\nabla \cap N = (\nabla_1 \cap N) \cup (\nabla_2 \cap N)$, so we deduce that the expressions for s and t above form honest global sections of $\mathcal{L}|_W$. Thus the map ϕ is well-defined away from the base locus of $\mathcal{L}|_W$ and the fibre of ϕ over $s = 0$ is as required. \square

It follows from the proof of Proposition 3.9 that the line bundle $\mathcal{O}_W(\hat{S})$ is just $\mathcal{O}_W(\sum_{\sigma \in (\Delta^\circ \setminus \nabla) \cap N} D_\sigma)$. Since W is an anticanonical hypersurface, the intersection of a divisor D_σ with W is empty if and only if σ lies in the relative interior of a facet of Δ° . If σ is in the interior of a codimension 2 face of Δ° , then a calculation analogous to that performed in [Roh04, §3.3] shows that $D_\rho \cap W$ has $1 + \ell^*(\Gamma(\sigma))\ell^*(\Gamma(\sigma)^\vee)$ irreducible components, where $\Gamma(\sigma)$ is the smallest face of Δ° containing σ , $\Gamma(\sigma)^\vee$ is the face of Δ made up of points ρ satisfying $\langle \sigma, \rho \rangle = -1$, and $\ell^*(\Gamma)$ denotes the number of lattice points in the relative interior of Γ . Finally, if σ lies in a codimension ≥ 3 face of Δ° , then $D_\rho \cap W$ is irreducible for generic W .

PROPOSITION 3.10. *If σ is in $(\Delta^\circ \setminus \nabla) \cap N$, then $D_\rho \cap W$ has a single irreducible component. Therefore, the member of the pencil \mathcal{P} corresponding to $[s : t] = [0 : 1]$ has*

$$\#(\Delta^\circ \setminus \nabla) \cap N$$

irreducible components.

PROOF. First, if σ is contained on the relative interior of a facet of Δ° , then $\Gamma(\sigma)^\vee$ is a single vertex η of Δ . Without loss of generality, we can assume that $\eta \in \Delta_1$. Therefore, $\langle \sigma, \rho \rangle \geq -1$ for all points $\rho \in \Delta$ and $\langle \sigma, \rho \rangle = -1$ if and only if

$\rho = \eta$, so, by definition, σ is in ∇_1 . Since, by [BB96, Proposition 6.3], all points of $\nabla \cap N$ are either in ∇_1 or ∇_2 , it follows that no point of $(\Delta^\circ \setminus \nabla) \cap N$ is in the interior of a facet of Δ° . Thus for any point $\sigma \in (\Delta^\circ \setminus \nabla) \cap N$, the intersection $D_\sigma \cap W$ is nonempty.

It just remains to treat the case where σ lies in a codimension 2 face of Δ° . Since σ lies in $(\Delta^\circ \setminus \nabla) \cap N$, by definition there must be some $\rho_1 \in \Delta_1$ and $\rho_2 \in \Delta_2$ so that $\langle \rho_1, \sigma \rangle = \langle \rho_2, \sigma \rangle = -1$. Therefore, $\Gamma(\sigma)^\vee$ contains points in both Δ_1 and Δ_2 , so is a face of neither. Given this, [BB96, Proposition 6.3] implies that $\Gamma(\sigma)^\vee$ does not contain any points in its relative interior. So $\ell^*(\Gamma(\sigma)^\vee) = 0$ and hence $D_\sigma \cap W$ has a single irreducible component. \square

REMARK 3.11. For W of arbitrary dimension, the same proof works to find the number of components of the member of the pencil \mathcal{P} corresponding to $[s : t] = [0 : 1]$. However, if $\dim(W) \geq 4$ we do not know whether this may be interpreted as a count of components of a singular fibre in a fibration on some birational model \hat{W} of W (see Remark 3.8).

Next we show that this number also has meaning with respect to the mirror Calabi-Yau variety V and its degeneration to the union of X_1 and X_2 .

PROPOSITION 3.12. *If $\dim(\nabla_1) = \dim(\nabla_2) = 4$, then $V \cap X_1 \cap X_2$ is an irreducible curve C of genus*

$$g(C) = \#(\Delta^\circ \setminus \nabla) \cap N - 1.$$

PROOF. By construction, C is a complete intersection of sections of the line bundles $\omega_{X_\Delta}^{-1}$ (which determines V), $\mathcal{L}_1 = \mathcal{O}_{X_\Delta}(\sum_{\rho \in \Delta_1 \cap N} D_\rho)$ (which determines X_1), and $\mathcal{L}_2 = \mathcal{O}_{X_\Delta}(\sum_{\rho \in \Delta_2 \cap N} D_\rho)$ (which determines X_2). The Koszul complex resolving \mathcal{O}_C is thus given by

$$\omega_{X_\Delta}^2 \rightarrow (\mathcal{L}_1^{-1} \otimes \omega_{X_\Delta}) \oplus (\mathcal{L}_2^{-1} \otimes \omega_{X_\Delta}) \oplus \omega_{X_\Delta} \rightarrow \mathcal{L}_1^{-1} \oplus \mathcal{L}_2^{-1} \oplus \omega_{X_\Delta} \rightarrow \mathcal{O}_{X_\Delta}$$

The corresponding second spectral sequence converges to $H^i(C, \mathcal{O}_C[3])$, so

$$\bigoplus_{p+q=i+3} {}''E_\infty^{p,q} \cong H^i(C, \mathcal{O}_C).$$

The relevant portion of ${}''E_1^{p,q}$ is given by

$$\begin{array}{ccccccc} H^4(\omega_{X_\Delta}^2) & \rightarrow & H^4(\mathcal{L}_1^{-1} \otimes \omega_{X_\Delta}) \oplus H^4(\mathcal{L}_2^{-1} \otimes \omega_{X_\Delta}) \oplus H^4(\omega_{X_\Delta}) & \rightarrow & \mathbb{C} & \rightarrow & 0 \\ 0 & \rightarrow & 0 & \rightarrow & 0 & \rightarrow & 0 \\ 0 & \rightarrow & 0 & \rightarrow & 0 & \rightarrow & 0 \\ 0 & \rightarrow & 0 & \rightarrow & 0 & \rightarrow & 0 \\ 0 & \rightarrow & 0 & \rightarrow & 0 & \rightarrow & \mathbb{C} \end{array}$$

Now, by [BB96, Theorem 2.5], we know that

$$\begin{aligned} h^4(\omega_{X_\Delta}^2) &= \ell^*(2\Delta^\circ), \\ h^4(\omega_{X_\Delta}) &= 1 \\ h^4(\mathcal{L}_1^{-1} \otimes \omega_{X_\Delta}) &= \ell^*(\nabla_1 + \Delta^\circ), \\ h^4(\mathcal{L}_2^{-1} \otimes \omega_{X_\Delta}) &= \ell^*(\nabla_2 + \Delta^\circ). \end{aligned}$$

It is not then hard to see that this spectral sequence degenerates at the ${}''E_2$ term and $h^0(\mathcal{O}_C) = 1$, hence C is irreducible. Since $h^i(\mathcal{O}_C) = 0$ for $i > 1$, we have that

the top row of $E_1^{p,q}$ above is exact except at the left-most term. Thus we can compute that

$$g(C) = \ell^*(2\Delta^\circ) - (\ell^*(\nabla_1 + \Delta^\circ) + \ell^*(\nabla_2 + \Delta^\circ)).$$

It remains to show that this is precisely the number of points in $(\Delta^\circ \setminus \nabla) \cap N$. For this we need a small lemma.

LEMMA 3.13. *If Q is either ∇_i or Δ° , the number $\ell^*(Q + \Delta^\circ)$ is equal to $\ell(Q)$, where $\ell(Q)$ denotes the number of lattice points in Q .*

PROOF. The polytope Δ° is defined by the inequalities $\langle \sigma, \rho \rangle \geq -1$ for all points $\rho \in \Delta$. Similarly, ∇_1 is defined by the inequalities $\langle \sigma, \rho \rangle \geq -1$ for all points $\rho \in \Delta_1$ and $\langle \sigma, \rho \rangle \geq 0$ for all points $\rho \in \Delta_2$. We shall prove the lemma for $Q = \nabla_1$; the other cases are analogous.

Now, the polytope $\nabla_1 + \Delta^\circ$ is defined by the inequalities $\langle \sigma, \rho \rangle \geq -2$ for $\rho \in \Delta_1$ and $\langle \sigma, \rho \rangle \geq -1$ for $\rho \in \Delta_2$. Therefore a point in the interior of $\nabla_1 + \Delta^\circ$ satisfies these inequalities strictly, and thus any lattice point in the interior of $(\nabla_1 + \Delta^\circ)$ has $\langle \sigma, \rho \rangle \geq 0$ for all $\rho \in \Delta_2$ and $\langle \sigma, \rho \rangle \geq -1$ for all $\rho \in \Delta_1$. But this is just the set of all lattice points in ∇_1 . \square

From this lemma, we see that

$$\ell^*(2\Delta^\circ) - (\ell^*(\nabla_1 + \Delta^\circ) + \ell^*(\nabla_2 + \Delta^\circ)) = \ell(\Delta^\circ) - \ell(\nabla_1) - \ell(\nabla_2).$$

Moreover, [BB96, Proposition 6.3] shows that all lattice points of ∇ are lattice points of either ∇_1 or ∇_2 , so this is equal to

$$\ell(\Delta^\circ) - \ell(\nabla) + 1 = \#(\Delta^\circ \setminus \nabla) \cap N + 1;$$

here the extra $(+1)$ term corresponds to the fact that we have over-counted the origin, which is the intersection of ∇_1 and ∇_2 . This completes the proof of Proposition 3.12. \square

REMARK 3.14. A very minor modification of this proof shows that, in the case where V has dimension $d \geq 3$, we have $h^{d-2,0}(V \cap X_1 \cap X_2) = \#(\Delta^\circ \setminus \nabla) \cap N - 1$. If $\dim(V) = 2$, then $h^0(V \cap X_1 \cap X_2) = \#(\Delta^\circ \setminus \nabla) \cap N$.

Putting everything together, we obtain the following theorem.

THEOREM 3.15. *If $\dim(V) = \dim(W) = d \geq 3$, then the member of the pencil of hypersurfaces \mathcal{P} corresponding to $[s : t] = [0 : 1]$ has exactly $h^{d-2,0}(V \cap X_1 \cap X_2) + 1$ components. If $\dim(V) = \dim(W) = 2$, then $V \cap X_1 \cap X_2$ is a set of points and the member of the pencil of hypersurfaces \mathcal{P} corresponding to $[s : t] = [0 : 1]$ has exactly $\#(V \cap X_1 \cap X_2)$ components.*

Next we analyze the rest of the members of the pencil \mathcal{P} on W . Our goal is to show that the members corresponding to $[1 : 0]$ and $[1 : 1]$ are essentially the singular fibres of the LG models of X_1 and X_2 . Thus there is a very real sense in which the pencil \mathcal{P} on W is collecting information about the LG models of X_1 and X_2 .

First, however, we describe how these LG models are constructed. In [Har16], it is shown that the naïve compactification of Givental's [Giv98] Landau-Ginzburg model for a complete intersection X in a toric variety X_Δ is smooth if X has dimension less than or equal to 3, and otherwise has only mild singularities.

This compactification is defined as follows. Assume that we have a polytope Δ and a nef partition Δ_1, Δ_2 of Δ , so that Δ_1 and Δ_2 contain no interior points. In this setting, a general enough global section of the line bundle \mathcal{L}_1 associated to Δ_1 determines a quasi-Fano hypersurface X in X_Δ . The compactified version of Givental’s LG model for X is then the complete intersection $Y \subset X_\nabla \times \mathbb{A}^1$ cut out by the equations

$$\sum_{\rho \in \Delta_1 \cap M} a_\rho \prod_{\sigma \in \nabla \cap N} z_\sigma^{\langle \sigma, \rho \rangle - \sigma_{\min}^1} = 0$$

$$ta_0 \prod_{\sigma \in \nabla_2 \cap N \setminus 0_N} z_\sigma - \sum_{\rho \in \Delta_2 \cap M \setminus 0_M} a_\rho \prod_{\sigma \in \partial \nabla \cap N} z_\sigma^{\langle \sigma, \rho \rangle - \sigma_{\min}^2} = 0$$

where a_ρ are complex constants, t is the coordinate on \mathbb{A}^1 , and σ_{\min}^i is -1 if σ is in ∇_i and 0 otherwise. The superpotential w is just projection of this complete intersection onto \mathbb{A}^1 . It is shown in [Har16] that (Y, w) has the expected properties for an LG model of X .

With this in place, we find:

THEOREM 3.16. *The members of the pencil \mathcal{P} corresponding to $[s : t] = [1 : 1]$ and $[1 : 0]$ are birational to the fibres over 0 of the LG models (Y_1, w_1) and (Y_2, w_2) of X_1 and X_2 respectively. In fact, for any choice of W and hypersurface $\mathcal{P}([1 : t])$ with $t \in \mathbb{C}$, there is a choice of LG model (Y, w) of either X_1 or X_2 so that $\mathcal{P}([1 : t])$ is birational to a fibre of (Y, w) .*

PROOF. Recall from Section 3.2 that we have an expression for a birational model of W as the complete intersection in $\mathbb{P}^1[s, t] \times X_\Delta^\circ$ given by the vanishing of

$$f_1 := s \sum_{\rho \in \Delta_1 \cap M \setminus 0_M} a_\rho \prod_{\sigma \in \partial \Delta^\circ \cap N} z_\sigma^{\langle \sigma, \rho \rangle + 1} - ta_0 \prod_{\sigma \in \partial \Delta^\circ \cap N} z_\sigma$$

$$f_2 := s \sum_{\rho \in \Delta_2 \cap M \setminus 0_M} a_\rho \prod_{\sigma \in \partial \Delta^\circ \cap N} z_\sigma^{\langle \sigma, \rho \rangle + 1} - (s - t)a_0 \prod_{\sigma \in \partial \Delta^\circ \cap N} z_\sigma.$$

Note that f_1 has a factor of $\prod_{\sigma \in \nabla_2 \cap N \setminus 0_N} z_\rho$ by the definition of ∇_2 , and an analogous statement holds for f_2 , by the definition of ∇_2 .

Now if we let $[s : t] = [1 : 1]$, then we obtain the complete intersection of

$$f_1|_{[1:1]} = \sum_{\rho \in \Delta_1 \cap M} a_\rho \prod_{\sigma \in \partial \Delta^\circ \cap N} z_\sigma^{\langle \sigma, \rho \rangle + 1}$$

$$f_2|_{[1:1]} = \sum_{\rho \in \Delta_2 \cap M \setminus 0_M} a_\rho \prod_{\sigma \in \partial \Delta^\circ \cap N} z_\sigma^{\langle \sigma, \rho \rangle + 1}.$$

Note that this is precisely the complete intersection determining the fibre over 0 of the LG model of X_1 , except compactified in X_{Δ° instead of X_∇ .

To compare these compactifications, define a birational map φ from X_{Δ° to X_∇ , which sends z_σ to z_σ if $\sigma \in \partial \nabla \cap N$. The restriction of φ to the complement of the torus invariant loci of codimension ≥ 2 simply has the effect of removing those codimension 1 tori corresponding to points $\sigma \in (\Delta^\circ \setminus \nabla) \cap N$. Thus we see that φ induces a birational map between $\mathcal{P}([1 : 1])$ and the fibre over 0 of (Y_1, w_1) if no components of $\mathcal{P}([1 : 1])$ are contained in torus invariant loci of X_{Δ° of codimension ≥ 2 , and no component of $\mathcal{P}([1 : 1])$ is contained in a divisor D_σ for $\sigma \in (\Delta^\circ \setminus \nabla) \cap N$.

The first of these two claims is trivial: since each component of $\mathcal{P}([1 : 1])$ is of codimension 2 in X_{Δ° , it is contained in a codimension ≥ 2 torus invariant subvariety of X_{Δ° if and only if it is the closure of such a torus invariant subvariety. Since W contains no torus invariant subvarieties of X_{Δ° , this cannot happen. The second claim follows from the fact that if $\sigma \in (\Delta^\circ \setminus \nabla) \cap N$, then $D_\sigma \cap W$ is in $\mathcal{P}([0 : 1])$. Thus $\mathcal{P}([1 : 1])$ can only intersect D_σ in at most a codimension 2 subvariety of X_{Δ° .

An identical argument suffices to show that $\mathcal{P}([1 : 0])$ is birational to the fibre over 0 of (Y_2, \mathbf{w}_2) , and in fact this shows that for any $p \in \mathbb{P}^1 \setminus \{[0 : 1]\}$, for a generic choice of W there is a choice of $i \in \{1, 2\}$ and an LG model (Y_i, \mathbf{w}_i) so that $\mathcal{P}(p)$ is birational to a fibre of (Y_i, \mathbf{w}_i) . \square

As a philosophical remark, this proves that all of the interesting data surrounding the pencil \mathcal{P} on W is related to either the LG models of X_1 and X_2 , or the variety $V \cap X_1 \cap X_2$. Indeed, this is the same data as was used to determine the Tyurin degeneration of V : X_1 and X_2 were the quasi-Fano hypersurfaces in X_Δ , and $V \cap X_1 \cap X_2$ was the locus that needed to be blown up in X_1 to obtain a smooth degeneration.

In the case where both X_1 and X_2 are pullbacks to X_Δ of ample hypersurfaces in \mathbb{P}_Δ along the mpcp resolution map, then we can say even more using the following theorem, proved in [Har16].

THEOREM 3.17. [Har16] *With notation as above, suppose in addition that X_i is the pull-back to X_Δ of an ample hypersurface in \mathbb{P}_Δ and that $\dim \Delta = d \geq 4$. Let (Y_i, \mathbf{w}_i) denote the LG model of X_i and let ρ_0 denote the number of components in its singular fibre over 0. Then*

$$\rho_0 = h^{d-2,1}(X_i) + 1.$$

From this, we immediately obtain:

COROLLARY 3.18. *Suppose that $\dim(V) = \dim(W) = 3$ and let ρ_p be the number of irreducible components in the member of the pencil \mathcal{P} corresponding to $p \in \mathbb{P}^1$. If X_1 and X_2 are pullbacks of ample hypersurfaces in \mathbb{P}_Δ , then*

- $\rho_{[1:0]} = h^{2,1}(X_1) + 1,$
- $\rho_{[1:1]} = h^{2,1}(X_2) + 1,$
- $\rho_{[0:1]} = h^{1,0}(C) + 1.$

REMARK 3.19. Note that the same result is true for \hat{W} , as the birational transformation from W to \hat{W} is an isomorphism in codimension 1. Thus the preceding corollary can be interpreted as a count of components in singular fibres of the K3 surface fibration $\pi: \hat{W} \rightarrow \mathbb{P}^1$ on \hat{W} .

Now let ℓ be the rank of the image of the restriction map $H^2(\hat{W}, \mathbb{C}) \rightarrow H^2(S, \mathbb{C})$, for S a smooth fibre of π . Using [DHNT16, Lemma 3.2], one can easily show that

$$h^{1,1}(\hat{W}) = \sum_{p \in \mathbb{P}^1} (\rho_p - 1) + \ell + 1$$

Moreover, by [Bat94] we see that $h^{2,1}(V) = h^{1,1}(W) = h^{1,1}(\hat{W})$. So, noting that $h^{2,1}(\hat{X}_1) = h^{2,1}(X_1) + g(C)$ (see, for instance, [Voi07, Theorem 7.31]),

Theorem 2.5 gives

$$\sum_{p \in \mathbb{P}^1 \setminus \{[1:0], [0:1], [1:1]\}} (\rho_p - 1) + \ell + k = 20.$$

This implies that if Dolgachev-Nikulin mirror symmetry does not hold (in a precise sense) for the K3 surfaces associated to the nef partition Δ_1, Δ_2 and their Batyrev-Borisov duals, then this failure is seen by the fibres of the fibration $\pi: \hat{W} \rightarrow \mathbb{P}^1$ away from the points in the set $\{[1:0], [0:1], [1:1]\}$.

Finally, one may ask whether an analogue of Corollary 3.18 holds when V and W are K3 surfaces. The difficulty here is in proving an analogue of Theorem 3.17: for subtle combinatorial reasons, the proof given in [Har16] does not easily generalize to the K3 surface case. However, we expect the following conjecture to hold in this case:

CONJECTURE 3.20. *Suppose that $\dim(V) = \dim(W) = 2$ and let ρ_p be the number of irreducible components in the fibre of the elliptic fibration $\pi: W \rightarrow \mathbb{P}^1$ over $p \in \mathbb{P}^1$. If X_1 and X_2 are pullbacks of ample hypersurfaces in \mathbb{P}_Δ , then*

- $\rho_{[1:0]} = h^{1,1}(X_1) - h^{1,1}(X_\Delta) + 1,$
- $\rho_{[1:1]} = h^{1,1}(X_2) - h^{1,1}(X_\Delta) + 1,$
- $\rho_{[0:1]}$ is the number of points in $V \cap X_1 \cap X_2$ and the corresponding singular fibre is semistable (Kodaira type I_n), and
- all other fibres of π are irreducible.

We will illustrate this conjecture with an example.

EXAMPLE 3.21 (Anticanonical hypersurfaces in $(\mathbb{P}^1)^3$). Let us take V to be an anticanonical hypersurface in $(\mathbb{P}^1)^3$. This is a K3 surface with Picard lattice of rank 3, isomorphic to the lattice with Gram matrix

$$\begin{pmatrix} 0 & 2 & 2 \\ 2 & 0 & 2 \\ 2 & 2 & 0 \end{pmatrix}.$$

There is a Tyurin degeneration of V to a union of two $(1, 1, 1)$ divisors X_1, X_2 in $(\mathbb{P}^1)^3$. The intersection $V \cap X_1 \cap X_2$ is 12 points. On the other side, we see that there is an elliptic fibration on the mirror dual K3 surface W , which has three reducible fibres of types I_{12}, I_2 and I_2 .

The polytope Δ determining $(\mathbb{P}^1)^3$ has vertices $\sigma_0, \dots, \sigma_5$ given by the columns of the matrix

$$\begin{pmatrix} 1 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & -1 \end{pmatrix}$$

The appropriate nef partition is $E_1 = \{\sigma_0, \sigma_2, \sigma_4\}$ and $E_2 = \{\sigma_1, \sigma_3, \sigma_5\}$, which has dual nef partition with

$$\begin{aligned} \nabla_1 &= \text{Conv}(\{(1, 0, 0), (0, 1, 0), (0, 0, 1), (1, 1, 1), (1, 1, 0), (0, 1, 1), (0, 1, 1), (0, 0, 0)\}) \\ \nabla_2 &= -\nabla_1. \end{aligned}$$

We draw the polytopes Δ°, ∇_1 and ∇_2 in Figure 2. The leftmost picture in Figure 2 is just the polytope Δ° , the middle picture denotes ∇_1 and ∇_2 and the picture on the right shows $(\Delta^\circ \setminus \nabla) \cap N$. It is clear from the description of the fibre over $[0:1]$ that it is actually semi-stable, so it follows from Kodaira’s classification




FIGURE 2. Polytopes related to Example 3.21

of singular fibres of elliptic fibrations that the resulting fibre is necessarily of type I_{12} . The same cannot be said for the fibres over $[1 : 0]$ and $[1 : 1]$, which have two components each, since it is not necessarily true that these fibres have normal crossings. Kodaira's classification can only be used to determine that these fibres are either of type I_2 or of type III.

4. Dolgachev-Nikulin mirror symmetry

Next we'll turn our attention to K3 surfaces. As noted in the previous section, in the setting of Batyrev-Borisov mirror symmetry, a refinement of the nef partition defining our K3 surface V gives rise to both a Tyurin degeneration of V and an elliptic fibration on the mirror K3 surface W . In the K3 surface case this appears to be part of a wider correspondence, which seems to have first been noticed by Dolgachev [Dol96], between Type II degenerations (of which Tyurin degenerations are an example) and elliptic fibrations on the Dolgachev-Nikulin mirror.

We begin by giving a precise statement of this correspondence. Suppose we have a (pseudo-ample) L -polarized K3 surface V , for some lattice L . To define the Dolgachev-Nikulin mirror of V , we first fix a primitive isotropic vector f in the orthogonal complement L^\perp of L in the K3 lattice $\Lambda_{K3} \cong H^{\oplus 3} \oplus E_8^{\oplus 2}$ (where H denotes the hyperbolic plane lattice and E_8 is the negative definite root lattice E_8). With this in place, the Dolgachev-Nikulin mirror W of V is defined to be an \check{L} -polarized K3 surface, where

$$\check{L} := (\mathbb{Z}f)^\perp_{L^\perp} / \mathbb{Z}f.$$

Note that this depends upon the choice of isotropic vector f .

By the discussion in [Dol96, Section 6] (see also [Sca87, Section 2.1]), fixing f is equivalent to fixing a 0-dimensional cusp (Type III point) in the Baily-Borel compactification \overline{D}_L of the period domain D_L of L -polarized K3 surfaces. Call this cusp P .

Then we have the following result, which is essentially contained in [Dol96, Remark 7.11]:

PROPOSITION 4.1. *With notation as above, there is a bijective correspondence between 1-dimensional cusps in \overline{D}_L that pass through P , and primitive isotropic vectors $e \in \check{L}$.*

REMARK 4.2. Since 1-dimensional cusps in \overline{D}_L correspond to Type II degenerations of V and isotropic vectors $e \in \check{L}$ correspond to elliptic fibrations on W , this gives rise to a correspondence between Type II degenerations of V and elliptic fibrations on W (up to automorphism).

PROOF OF PROPOSITION 4.1. Suppose first that we have a 1-dimensional cusp $C \subset \overline{D}_L$ that passes through P . By [Sca87, Section 2.1], such cusps are in bijection with rank two primitive isotropic sublattices of L^\perp that contain the vector f . So C gives rise to a sublattice E of L^\perp and, since E is a primitive sublattice of $(\mathbb{Z}f)_{L^\perp}^\perp$ and E contains f , we see that $E \cap \check{L}$ is a primitive isotropic sublattice of \check{L} of rank 1. Let e denote a generator of this sublattice; then e is a primitive isotropic vector in \check{L} .

Conversely, suppose we have a primitive isotropic vector $e \in \check{L}$. By definition of \check{L} , we have $\langle e, f \rangle = 0$. So the lattice E spanned by e and f is a rank two primitive isotropic sublattice of L^\perp . But this gives rise to a 1-dimensional cusp passing through P . \square

We will now illustrate this correspondence in some explicit examples, which will provide a glimpse of some possible deeper structure.

4.1. H-polarized K3's. We begin by looking at the Type II degenerations of H -polarized K3's. An H -polarized K3 surface may be constructed as an anticanonical hypersurface in an mpcp resolution of the weighted projective space $\mathbb{WP}(1, 1, 4, 6)$. However, the defining polytope of this weighted projective space does not admit any nef partitions, so we cannot apply the theory of Section 3 to study it.

Instead, we will try a different approach to comparing Type II degenerations and elliptic fibrations on the mirror. Note first that an H -polarized K3 surface V naturally corresponds to a double cover of the Hirzebruch surface \mathbb{F}_4 , ramified over a divisor in the linear system $|4s + 12f|$, where s is the class of the (-4) -section and f is the class of a fibre (when there is no risk of confusion, we will always denote the $(-n)$ -section in a Hirzebruch surface \mathbb{F}_n by s and a fibre of the ruling by f). So H -polarized K3 surfaces are equivalent to pairs (\mathbb{F}_4, B) , where $B \in |4s + 12f|$ is the branch divisor.

The moduli space of such pairs admits a *KSBA compactification* (see [Ale96a, Ale96b, Ale06, KSB88]), which has been studied in detail by Brunyate [Bru15]. This compactification admits a birational morphism to the usual Baily-Borel compactification of the moduli space of H -polarized K3 surfaces. The Type II degenerations occur along two boundary components in this KSBA compactification, which map to the two 1-dimensional cusps in the Baily-Borel compactification. These two kinds of Type II degenerations may be described as follows:

- (1) Degenerate V to a union $X_1^1 \cup_Z X_2^1$ defined as follows. X_1^1 is a double cover of \mathbb{F}_4 ramified over a smooth divisor in the linear system $|2s + 12f|$, and $Z \subset X_1^1$ is an elliptic curve given by the pull-back of the (-4) -section. X_2^1 is also a double cover of \mathbb{F}_4 , this time ramified over a smooth divisor in the linear system $|s + 4f|$, and $Z \subset X_2^1$ is the pull-back of a second smooth divisor in the linear system $|s + 4f|$.

Riemann-Roch easily yields that there is a 17-dimensional space of deformations of (X_1^1, Z) preserving the double covering structure, and a 1-dimensional space of deformations of (X_2^1, Z) . To glue these components together along Z , we need to ensure that the elliptic curves Z in each component are isomorphic; this imposes a single gluing condition. The total configuration therefore has $17 + 1 - 1 = 17$ moduli. It is easy to see that any variety $X_1^1 \cup_Z X_2^1$ defined in this way is d -semistable (in the

sense of [Fri83]), so admits a smoothing to a K3 surface by [Fri83, Theorem 5.10]. Thus, such degenerations lie along a 17-dimensional boundary component (i.e. a boundary divisor) in the KSBA moduli space.

- (2) Degenerate V to a union $X_1^2 \cup_Z X_2^2$, where both X_1^2 and X_2^2 are double covers of \mathbb{F}_2 ramified over smooth divisors in the linear system $|4s + 6f|$ and Z is the pull-back of a fibre of the ruling on \mathbb{F}_2 . Riemann-Roch yields that each component (X_i^2, Z) has 9 deformations that preserve the double covering structure and, as before, there is a single gluing condition along Z . The total configuration therefore has $9 + 9 - 1 = 17$ moduli and is d -semistable, so gives another boundary divisor of the moduli space.

Now we look at the mirror. Up to isometry, there are two 1-dimensional cusps in \overline{D}_H , meeting in a unique 0-dimensional cusp. So we only have one choice of mirror W : a K3 surface polarized by the lattice $M := H \oplus E_8 \oplus E_8$.

Now we match the Type II degenerations above with elliptic fibrations on the mirror. From [CD07, Section 3.3], we know that an M -polarized K3 surface admits two elliptic fibrations:

- (1) The alternate fibration, which has an I_{12}^* and six I_1 's. Note that the I_{12}^* fibre has 17 components, corresponding to the 17 moduli of the component X_1^1 in degeneration (1) above.
- (2) The standard fibration, which has two II^* 's and four I_1 's. Note that each II^* fibre has 9 components, corresponding to the 9 moduli of each of the components X_i^2 in the Type II degeneration (2) above.

4.2. K3 surfaces of degree two. Now consider K3 surfaces of degree two (i.e. polarized by the rank one lattice $\langle 2 \rangle$). A K3 surface V of degree 2 may be constructed as an anticanonical hypersurface in the weighted projective space $\mathbb{WP}(1, 1, 1, 3)$. This toric variety is determined by the polytope $\Delta \subset M_{\mathbb{R}} \cong \mathbb{R}^3$ with vertices given by the columns $\sigma_1, \sigma_2, \sigma_3, \sigma_4$ of the matrix

$$\begin{pmatrix} 1 & 0 & 0 & -1 \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & -3 \end{pmatrix}.$$

Up to automorphism, there is only one bipartite nef partition of this polytope, given by $E_1 = \{\sigma_1, \sigma_2, \sigma_4\}$ and $E_2 = \{\sigma_3\}$. The dual nef partition in $N_{\mathbb{R}}$ has

$$\nabla_1 = \text{Conv}(-e_1^* - e_2^*, -e_1^* - e_2^* + e_3^*, -e_1^* + 2e_2^*, 2e_1^* - e_2^*)$$

$$\nabla_2 = \text{Conv}(0_N, -e_3^*, 3e_1^* - e_3^*, 3e_2^* - e_3^*)$$

where e_1, e_2 and e_3 are basis vectors of $M_{\mathbb{R}}$ and e_i^* are their duals.

As in Example 3.21, we may look at the polytopes ∇_1 and ∇_2 in comparison to Δ° and note that the integral points in $\Delta^\circ \setminus \nabla$ form a cycle of length 18. Thus the fibre over $[0 : 1]$ in the elliptic fibration on the Batyrev dual W described in Section 3.2 is of type I_{18} . One can also check, using techniques described in [Har16], that the fibres over $[1 : 0]$ and $[1 : 1]$ are irreducible.

Finally, in the degree two case it is well known that the Batyrev dual family of K3 surfaces is actually Dolgachev-Nikulin dual (see, for instance, [Roh04]), so a generic Batyrev dual K3 surface W has Picard lattice $M_2 := H \oplus E_8 \oplus E_8 \oplus A_1$. The elliptic fibrations on such an M_2 -polarized K3 surface were computed by Dolgachev [Dol96, Remark 7.11]; we see that one of them has an I_{18} fibre and six I_1 's, as expected.

However, it is known (see, for instance, [Sca87, Section 6]) that the Baily-Borel compactification of the moduli space of K3 surfaces of degree two has four 1-dimensional cusps, corresponding to four types of Type II degenerations, yet the example above only gives one. We can analyse the others using the same techniques that we used in the H -polarized case above.

Indeed, it is well known that a K3 surface of degree two V naturally corresponds to a double cover of \mathbb{P}^2 ramified over a smooth sextic curve. The moduli space of K3 surfaces of degree two is therefore the same as the moduli space of pairs (\mathbb{P}^2, B) where B is a sextic curve.

Alexeev and Thompson have studied a KSBA compactification [AT16] for the moduli space of such pairs. The Type II degenerations occur along four boundary components, which map to the four cusps in the Baily-Borel compactification. The four corresponding kinds of Type II degenerations may be described as follows:

- (1) Degenerate V to a union $X_1^1 \cup X_2^1$, defined as follows. X_1^1 is a double cover of \mathbb{F}_4 ramified over a smooth divisor in the linear system $|2s + 12f|$, and $Z \subset X_1^1$ is an elliptic curve given by the pull-back of the (-4) -section. X_2^1 is a double cover of \mathbb{P}^2 ramified over a smooth conic, and $Z \subset X_2^1$ is the pull-back of a second smooth conic. This degeneration is in many ways analogous to degeneration (1) in the H -polarized case.

In this case the pair (X_1^1, Z) admits 16 deformations preserving the double covering structure, and (X_2^1, Z) has two. As in the H -polarized case, there is a single gluing condition corresponding to choice of the elliptic curve Z , so the total configuration has $17 + 2 - 1 = 18$ moduli. It is easy to see that all such varieties are d -semistable, so they give rise to an 18-dimensional boundary component (i.e. a boundary divisor) in the KSBA moduli space.

- (2) Degenerate V to a union $X_1^2 \cup_Z X_2^2$, defined as follows. X_1^2 is a double cover of \mathbb{F}_1 ramified over a smooth divisor in the linear system $|2s + 6f|$ and $Z \subset X_1^2$ is the pull-back of the (-1) -section. X_2^2 is a double cover of \mathbb{P}^2 ramified over a smooth quartic and $Z \subset X_2^2$ is the pull-back of a line.

Counting deformations that preserve the double cover structure, we see that (X_1^2, Z) has 11 deformations and the (X_2^2, Z) component has 8. However, as usual there is a gluing condition along Z , so the total configuration therefore has $11 + 8 - 1 = 18$ moduli. Since such varieties are all d -semistable, they correspond to a boundary divisor in the moduli space.

- (3) The last two cases are more interesting. In the first, we degenerate V to a single component X_1^3 , defined to be the double cover of a cubic cone ramified over a smooth anticanonical divisor. X_1^3 thus contains two log canonical singularities, corresponding to the pull-backs of the vertex of the cone. This double cover has 18 moduli, and corresponds to a boundary divisor in the KSBA moduli space.

Interestingly, this degeneration may be seen to be equivalent to the degeneration given by the nef partition above. Indeed, if $X_1 \cup_Z X_2$ is the degeneration corresponding to this nef partition, we may obtain the degeneration X_1^3 by blowing up along Z , then contracting X_1 and X_2 . These contractions give rise to the two log canonical singularities present in X_1^3 .

- (4) The final case is most difficult. In [AT16], the corresponding degeneration of V is a union $X_1^4 \cup_Z X_2^4$, where each X_i^4 is a double cover of $\mathbb{F}_2^0 \cong \mathbb{W}\mathbb{P}(1, 1, 2)$ (obtained by taking a copy of \mathbb{F}_2 and contracting the (-2) -section) ramified over a divisor in the linear system $\mathcal{O}(6)$ and cyclically over the A_1 singularity, and Z is the pull-back of a divisor in the linear system $\mathcal{O}(1)$. This degeneration is in many ways analogous to degeneration (2) in the H -polarized case.

Counting moduli, we see that each component (X_i^4, Z) has 9 deformations that preserve the double covering structure, but there is, as always, one gluing condition along Z . We thus have $9 + 9 - 1 = 17$ deformations, which does not give the expected divisor on the boundary. However, there is also something wrong with this degeneration: one can easily check that it is not d -semistable, so does not appear as the central fibre in a smooth semistable degeneration of K3 surfaces.

There are two ways to resolve this. The first may be thought of as analogous to the blow-up of $V \cap X_1 \cap X_2$ in X_1 from Section 3: we simply blow up an arbitrary point on the image of Z in one copy of \mathbb{F}_2^0 . This corresponds to blowing up a pair of points on Z in X_1^4 , under the condition that the two points blown up are exchanged by the involution defining the double covering. After blowing up, $X_1^4 \cup_Z X_2^4$ becomes d -semistable and acquires an extra modulus, corresponding to the choice of point to blow up, making the corresponding boundary component into a divisor.

However, there is also a second way to resolve this problem. Inspired by (3), above, we could try introducing a third component between X_1^4 and X_2^4 ; this has the appeal of maintaining the symmetry of the degenerate fibre. Such a central component X_3^4 can be constructed as follows. Take a copy of \mathbb{P}^2 and blow up three points in general position, to get a (-1) -hexagon. Blow up one of the vertices of this hexagon again, then contract the two (-2) -curves that result. One obtains a surface that has two A_1 singularities, and the (-1) -hexagon becomes a pentagon whose sides have self-intersections $(-1, -1, -\frac{1}{2}, 0, -\frac{1}{2})$. The sum of the two (-1) -curves in this pentagon gives a ruling and X_3^4 is a double cover of this surface ramified over three of its fibres, as well as cyclically over both of the A_1 singularities. X_3^4 glues to X_1^4 and X_2^4 along two isomorphic elliptic curves Z_1 and Z_2 , which are the pull-backs of the two $(-\frac{1}{2})$ -curves. We thus obtain a degenerate fibre $X_1^4 \cup_{Z_1} X_3^4 \cup_{Z_2} X_2^4$.

Accounting for automorphisms, (X_3^4, Z_1, Z_2) has 2 deformations. So, with this component included, the total moduli count is $9+9+2-1-1 = 18$ (the two (-1) 's appear because there is a gluing condition associated to each double curve). Moreover, the fibre $X_1^4 \cup_{Z_1} X_3^4 \cup_{Z_2} X_2^4$ is d -semistable, so we get a boundary divisor in moduli.

Now we again match with elliptic fibrations on the mirror. These are computed by Dolgachev [Dol96, Remark 7.11] as:

- (1) An elliptic fibration which has one I_{12}^* , one I_2 , and four I_1 's. Note that the I_{12}^* fibre has 17 components, corresponding to the 17 moduli of (X_1^1, Z) in degeneration (1) above, and the I_2 fibre has two components, corresponding to the two moduli of (X_2^1, Z) .

- (2) An elliptic fibration with one fibre of type I_6^* , one fibre of type III^* , and three I_1 's. As above, the I_6^* fibre has 11 components, corresponding to the 11 moduli of (X_1^2, Z) in degeneration (4) above, and the III^* fibre has eight components, corresponding to the eight moduli of (X_2^2, Z) .
- (3) An elliptic fibration with one fibre of type I_{18} and six I_1 's. Once again, the I_{18} fibre has 18 components, corresponding to the 18 moduli of the single component X_3^1 in degeneration (3), above.
- (4) An elliptic fibration which has two II^* 's, one I_2 , and two I_1 's. Note that each II^* fibre has 9 components, corresponding to the 9 moduli of (X_1^4, Z) and (X_2^4, Z) in degeneration (2) above, and the I_2 fibre has two components. These two components can be thought of as corresponding to the two points on Z which are blown up to make $X_1 \cup_Z X_2$ d -semistable (c.f. Conjecture 3.20), or as corresponding to the two moduli of the “extra” central component (X_3^4, Z_1, Z_2) .

4.3. Discussion. These results are highly suggestive of the idea that the correspondence between Tyurin degenerations and elliptic fibrations on Batyrev dual K3 surfaces, explored in Section 3, may extend to a broader correspondence between Type II degenerations of an L -polarized K3 surface V and elliptic fibrations on its Dolgachev-Nikulin mirror W . Such a correspondence should have properties that generalize those described in Section 3; the aim of this section is to discuss the form that such properties may take.

Firstly, cases (3) and (4) of the considerations in Section 4.2 suggest that there may be some correspondence between the choice of model for the Type II degeneration of V and some properties of the elliptic fibration. Thinking about this in the context of the threefold philosophy outlined in Remark 2.7, one might conjecture the following.

Suppose first that we have a Type II degeneration of V to a configuration $X_1 \cup_Z X_2 \cup_Z \dots \cup_Z X_k$, where each X_i meets X_{i-1} and X_{i+1} . Let n_i be the number of deformations of (X_i, Z) (where Z denotes the double locus on X_i and may have more than one component) that preserve some notion of lattice polarization – this last condition is to ensure that the deformed Type II fibres still smooth to L -polarized K3 surfaces, and was arranged in the preceding examples by requiring that deformations preserve the double covering structure; it also accounts for the appearance of the $h^{1,1}(X_\Delta)$ term in Conjecture 3.20. Then the fibre dimension of the natural map from the moduli space of such pairs (X_i, Z) to the moduli space of elliptic curves Z is equal to $n_i - 1$.

Thus, noting that all elliptic double curves in the Type II degeneration are isomorphic, we see that such a Type II fibre should have $\sum_{i=1}^k (n_i - 1) + 1$ deformations. We first conjecture that this number should equal $19 - \ell$, where $\ell = \text{rank}(L)$. Then, since the moduli space of V (as an L -polarized K3 surface) is $(20 - \ell)$ -dimensional, such Type II degenerations should lie along codimension 1 loci in an appropriate compactification. We would thus get a decomposition of the $20 - \ell$ deformations of V into contributions $(n_i - 1)$ from deformations of each X_i , a contribution 1 from deformations of Z , and 1 for the codimension in the moduli space.

Now we look at the mirror picture. As proved in Proposition 4.1, the Type II degeneration of V given above should correspond to an elliptic fibration $\pi: W \rightarrow \mathbb{P}^1$ on the mirror W . We suggest that the decomposition $X_1 \cup_Z X_2 \cup_Z \dots \cup_Z X_k$ of V corresponds to a “slicing” of the \mathbb{P}^1 base of π , so that each X_i corresponds to

a slice S_i , as follows. S_1 is a disc, which is glued along its boundary to one of the boundaries of an annulus S_2 . The other boundary of the annulus S_2 is then glued to one of the boundaries of an annulus S_3 , and so on, until the remaining boundary of the annulus S_{k-1} is glued to a disc S_k .

The singular fibres of π should then be apportioned amongst the slices as follows. If ρ_p denotes the number of components in the fibre of π over p , then the slicing should satisfy

$$n_i - 1 = \sum_{p \in S_i} (\rho_p - 1).$$

This gives a decomposition of the Picard rank $(20 - \ell)$ of W into contributions $n_i - 1$ from the singular fibres lying on the slice S_i , a contribution 1 from the class of a section, and a contribution 1 from the class of a fibre.

Note that this is completely compatible with the Tyurin degeneration picture presented in Section 2 and the Batyrev-Borisov picture of Conjecture 3.20. Indeed, in the Tyurin degeneration picture the slicing has two pieces S_1 and S_2 , which are the LG models of the two components X_1 and X_2 of the degenerate fibre $X_1 \cup_{\mathbb{Z}} X_2$. Furthermore, in the setting of Conjecture 3.20 we also have a slicing into two pieces S_1 and S_2 , one of which contains the point $[1 : 0]$ and the other of which contains $[1 : 1]$. Which slice the point $[0 : 1]$ falls into depends upon the choice of blow-up of $V \cap X_1 \cap X_2$: if we blow up $V \cap X_1 \cap X_2$ in X_i (for $i \in \{1, 2\}$), then the corresponding I_n fibre appears in the slice S_i .

Finally, we describe how this slicing picture works in the H -polarized and degree 2 cases considered above. In the H -polarized case we have the following two possibilities, numbered compatibly with Section 4.1. The labelling of the slices is chosen so that the slice corresponding to the component X_i^j is labelled S_i^j .

- (1) \mathbb{P}^1 is sliced into two pieces S_1^1 and S_2^1 , such that S_1^1 contains the I_{12}^* fibre.
- (2) \mathbb{P}^1 is sliced into two pieces S_1^2 and S_2^2 , each of which contains a II^* fibre.

Moreover, in the degree two case we have the following four possibilities, numbered compatibly with Section 4.2.

- (1) \mathbb{P}^1 is sliced into two pieces S_1^1 and S_2^1 , such that S_1^1 contains the I_{12}^* fibre and S_2^1 contains the I_2 .
- (2) \mathbb{P}^1 is sliced into two pieces S_1^2 and S_2^2 , such that S_1^2 contains the I_6^* fibre and S_2^2 contains the III^* .
- (3) In this case we have two choices of slicing, corresponding to the two types of degeneration. For the degeneration given by the nef partition and Batyrev mirror symmetry, \mathbb{P}^1 is sliced into two pieces S_1 and S_2 , one of which contains the I_{18} fibre. For the KSBA degeneration, we have a “degenerate” slicing of \mathbb{P}^1 into a single piece, which contains all singular fibres.
- (4) In the final case we also have two choices of slicing, corresponding to the two types of degeneration. In the case where we blow up a pair of points, \mathbb{P}^1 is sliced into two pieces S_1^4 and S_2^4 , such that S_1^4 contains the II^* fibre and the I_2 fibre, and S_2^4 contains the other II^* . In the case where we have three components in the degeneration, \mathbb{P}^1 is sliced into three pieces S_1^4 , S_3^4 and S_2^4 , such that S_1^4 and S_2^4 are discs containing one II^* fibre each, and S_3^4 is an annulus containing the I_2 fibre.

5. Beyond Batyrev-Borisov mirror symmetry for threefolds

The aim of this section is to provide some evidence that the ideas presented in Section 2 also hold for threefolds outside the toric setting considered in Section 3. We begin by showing that classical mirror symmetry suggests a correspondence between Tyurin degenerations and K3 fibrations on mirror dual pairs of Calabi-Yau threefolds. This should be thought of as a threefold analogue of Proposition 4.1. Then we specialize our discussion to the case of threefolds fibred by mirror quartics, as studied in [DHNT16], and show that, in that setting, the correspondence predicted by classical mirror symmetry is consistent with the construction presented in Section 2.

5.1. Classical mirror symmetry for threefolds. Classical mirror symmetry predicts that if V and W are mirror dual Calabi-Yau threefolds, then there is a relation between monodromy operators acting on $H^3(V, \mathbb{Q})$ and divisors in the closure of the Kähler cone of W . We will briefly sketch some of the details of this relationship here, the interested reader may find more details in [CK99, Chapter 6].

Suppose that $\mathcal{V} \rightarrow (\Delta^*)^n$ is a family of Calabi-Yau threefolds over the punctured polydisc, with fibre $\mathcal{V}_t = V$ above some $t \in (\Delta^*)^n$. For each $i \in \{1, \dots, n\}$, let T_i be the unipotent monodromy operator acting on $H^3(V, \mathbb{Q})$ coming from the loop $(t_1, \dots, t_{i-1}, e^{2\pi i t}, t_{i+1}, \dots, t_n)$, where $(t_1, \dots, t_{i-1}, t_{i+1}, \dots, t_n)$ are fixed constants, and let $N_i = \log(T_i)$. The family \mathcal{V} is said to have *maximally unipotent monodromy* at $(0, \dots, 0)$ if

- (1) for any n -tuple (a_1, \dots, a_n) of positive integers, the weight filtration W_\bullet on $H^3(V, \mathbb{Q})$ induced by $\sum_{i=1}^n a_i N_i$ has $\dim W_0 = \dim W_1 = 1$ and $\dim W_2 = n + 1$, and
- (2) if g_0, \dots, g_n is a basis of W_2 chosen so that g_0 spans W_0 , and m_{ij} are defined by $N_i g_j = m_{ij} g_0$, then the matrix (m_{ij}) is invertible.

If \mathcal{V} has maximally unipotent monodromy, then mirror symmetry should produce a map which assigns to each N_i a divisor D_i in the closure of the Kähler cone of W . Moreover, there should be an identification under mirror symmetry which gives an isomorphism $H^{3-i,i}(V) \cong H^{i,i}(W)$, and hence an isomorphism $H^3(V, \mathbb{C}) \cong \bigoplus_{i=0}^3 H^{i,i}(W)$, so that the action of N_i on $H^3(V, \mathbb{C})$ agrees with the action of the cup product operator $J_i(-) = (-) \cup [-D_i]$ under this correspondence. Thus, for any n -tuple (a_1, \dots, a_n) of non-negative integers, the weight filtration on $H^3(V, \mathbb{C})$ induced by $N := \sum_{i=1}^n a_i N_i$ should be mirrored by the filtration on $\bigoplus_{i=0}^3 H^{i,i}(W)$ induced by $J := \sum_{i=1}^n a_i J_i$, and the limit Hodge decomposition should correspond to the decomposition $\bigoplus_{i=0}^3 H^{i,i}(W)$.

Now we specialize this discussion to the case of a Tyurin degeneration of Calabi-Yau threefolds $\mathcal{V} \rightarrow \Delta$. As in the previous sections, we write the central fibre of \mathcal{V} as $X_1 \cup_Z X_2$ and let V denote a general fibre. Let T be the monodromy operator acting on $H^3(V, \mathbb{Q})$ associated to a counterclockwise loop around 0. In order to apply the predictions of mirror symmetry, we assume that T may be identified with a loop $\prod_{i=1}^n T_i^{a_i}$ around a point of maximally unipotent monodromy in the complex moduli space of V , where T_i are as above and a_i are non-negative integers. Define $N := \log(T) = \sum_{i=1}^n a_i N_i$. We will use the Clemens-Schmid exact sequence associated to N to compute the limit mixed Hodge structure on $H^3(V)$, then see what this allows us to deduce about the mirror threefold W .

REMARK 5.1. We note that the Tyurin degeneration \mathcal{V} cannot have maximally unipotent monodromy, for purely topological reasons (see, for instance, [Mor84, Corollary 2]), so T must correspond to a loop around some positive-dimensional boundary component of the compactified complex moduli space of V . In particular, this implies that some of the a_i must be zero.

We begin by looking at the mixed Hodge structure on $H^3(\mathcal{V})$ given by Griffiths and Schmid [GS75, Section 4]. The weight filtration W_\bullet on $H^3(\mathcal{V}, \mathbb{Q})$ has

$$\text{Gr}_3^W = H^3(X_1, \mathbb{Q}) \oplus H^3(X_2, \mathbb{Q})$$

and, if r_1 and r_2 are the restriction maps $r_i: H^2(X_i, \mathbb{Q}) \rightarrow H^2(Z, \mathbb{Q})$, then

$$\text{Gr}_2^W = H^2(Z, \mathbb{Q}) / (\text{im}(r_1) + \text{im}(r_2)).$$

These weight graded pieces are then equipped with the appropriate Hodge filtrations. Define integers $u := \text{rank}(\text{Gr}_2^W) - 2$ and $v := \frac{1}{2} \text{rank}(\text{Gr}_3^W)$. Noting that K_{X_i} is anti-effective, so that $h^{3,0}(X_i) = 0$, we see that $v = h^{2,1}(X_1) + h^{2,1}(X_2)$.

The Clemens-Schmid exact sequence gives us an exact sequence of mixed Hodge structures

$$\dots \rightarrow H_5(\mathcal{V}) \rightarrow H^3(\mathcal{V}) \xrightarrow{i^*} H_{\text{lim}}^3(V) \xrightarrow{N} H_{\text{lim}}^3(V) \rightarrow H_3(\mathcal{V}) \rightarrow \dots$$

where i^* is the pull-back on cohomology induced by the inclusion $i: V \hookrightarrow \mathcal{V}$.

LEMMA 5.2. $H_5(\mathcal{V}) = 0$, so the map i^* is an injection.

PROOF. The Mayer-Vietoris sequence for $X_1 \cup_Z X_2$ gives

$$\dots \rightarrow H_5(X_1) \oplus H_5(X_2) \rightarrow H_5(\mathcal{V}) \rightarrow H_4(Z) \xrightarrow{\alpha} H_4(X_1) \oplus H_4(X_2) \rightarrow \dots,$$

where the map α is induced by the inclusions $Z \hookrightarrow X_i$.

Now, $H_5(X_1) \oplus H_5(X_2)$ vanishes by Poincaré duality and the assumption that $h^{0,1}(X_i) = 0$. Moreover, as Z is an effective anticanonical divisor in both X_1 and X_2 , the image of the class $[Z] \in H_4(Z)$ of Z under α is non-trivial. But $[Z]$ generates $H_4(Z)$, so α must be injective. Thus the sequence above gives $H_5(\mathcal{V}) = 0$. \square

Applying this lemma and some standard results on the Clemens-Schmid exact sequence (see, for instance, [Mor84]) we obtain the following limit mixed Hodge structure on $H^3(V)$

	Gr_F^3	Gr_F^2	Gr_F^1	Gr_F^0
Gr_4^M	\mathbb{C}	\mathbb{C}^u	\mathbb{C}	0
Gr_3^M	0	\mathbb{C}^v	\mathbb{C}^v	0
Gr_2^M	0	\mathbb{C}	\mathbb{C}^u	\mathbb{C}

where M_\bullet is the monodromy weight filtration induced by N and F^\bullet is the limit Hodge filtration.

Therefore, the divisor $D = \sum_{i=1}^n a_i D_i$ on W which corresponds to N under mirror symmetry should have

	$H^{0,0}(Y^\vee)$	$H^{1,1}(W)$	$H^{2,2}(W)$	$H^{3,3}(W)$
$\text{coimage}(J)$	\mathbb{C}	\mathbb{C}^u	\mathbb{C}	0
$\text{ker}(J)/\text{im}(J)$	0	\mathbb{C}^v	\mathbb{C}^v	0
$\text{im}(J)$	0	\mathbb{C}	\mathbb{C}^u	\mathbb{C}

where, as before, $J(-) = (-) \cup [-D]$ denotes the cup-product operator. In particular, we see that $J^2 = 0$. Since D is in the closure of the Kähler cone of W , results

of Oguiso [Ogu01, Example 2.3] show that mD is the class of a fibre in a fibration of W by K3 or abelian surfaces, for some positive integer m .

REMARK 5.3. Based on the ideas in the previous sections, we conjecture that mD will always be the class of fibre in a K3 fibration on W . Oguiso [Ogu01, Example 2.3] gives a simple criterion to test for this: mD defines a K3 fibration on W if and only if $c_2(W) \cdot D > 0$.

In light of this remark, we will assume throughout the remainder of this section that mD defines a K3 fibration on W . Then the calculation above also shows that the classes in $\text{Pic}(W)$ supported on fibres span a $v + 1$ dimensional subspace, where one of these classes is mD itself. Moreover, there is a rank u subspace of $\text{Pic}(W)$ with $J(\eta) \neq 0$ for each class $\eta \neq 0$ in this subspace. By the global invariant cycles theorem, classes in this second subspace come from monodromy invariant cycles on fibres of the K3-fibration on W . Thus the K3 surface fibration on W induced by mD is \check{L} -polarized (in the sense of [DHNT15, Definition 2.1]), for some lattice \check{L} of rank u .

Therefore we see that, if V admits a Tyurin degeneration to a union of threefolds $X_1 \cup_Z X_2$, and if restriction of divisors from X_1 and X_2 induces a lattice polarization of Z by a lattice L of rank $20 - u$, then we expect the mirror W to admit an \check{L} -polarized K3 surface fibration, for some lattice \check{L} of rank u . Moreover, the space of divisors in W that are supported on fibres of the fibration should have rank $v + 1 = h^{2,1}(Y_1) + h^{2,1}(Y_2) + 1$. Note that this is completely consistent with the predictions of Section 2.3.

5.2. Threefolds fibred by mirror quartics. Our next aim is to demonstrate how this works in a special case: that of threefolds fibred by mirror quartic K3 surfaces. As we will see, in this setting the predictions of classical mirror symmetry, described above, mesh perfectly with the construction presented in Section 2.

A detailed study of threefolds fibred by mirror quartic K3 surfaces was conducted in [DHNT16]. We begin by briefly recapping the main construction and results of that paper, before describing how it fits into our picture.

The goal of [DHNT16] is to answer the following question: let W be a Calabi-Yau threefold and assume that W admits a fibration over \mathbb{P}^1 by K3 surfaces, $\pi: W \rightarrow \mathbb{P}^1$. Assume that the general fibre of π is a K3 surface S with $\text{Pic}(S) = M_2$, for $M_2 := H \oplus E_8 \oplus E_8 \oplus \langle -4 \rangle$ (i.e. S is Dolgachev-Nikulin mirror to a quartic hypersurface in \mathbb{P}^3), and that the monodromy representation acts trivially on M_2 . Such a structure is called an M_2 -polarized K3 fibration on W . In [DHNT16] we attempted to classify Calabi-Yau threefolds admitting M_2 -polarized K3 fibrations.

In order to describe this classification, we start by taking a basic family, called \mathcal{X} in [DHNT16], which is a smooth resolution of the family of hypersurfaces

$$\{\lambda w^4 + xyz(x + y + z - w) = 0\} \subset \mathbb{P}^3,$$

for $\lambda \neq 1/256, 0$. This defines a smooth family of K3 surfaces over $\mathbb{P}^1 \setminus \{0, 1/256, \infty\}$. The classification of M_2 -polarized fibrations from [DHNT16] can then be stated as:

THEOREM 5.4. [DHNT16, Section 2] *If $\pi: W \rightarrow \mathbb{P}^1$ is an M_2 -polarized K3 fibration on a Calabi-Yau threefold W , then there is a map*

$$g: \mathbb{P}^1 \rightarrow \mathbb{P}^1$$

so that W is birational to $g^* \mathcal{X}$ where

$$\begin{array}{ccc}
 g^* \mathcal{X} & \longrightarrow & \mathcal{X} \\
 \downarrow & & \downarrow \\
 U & \xrightarrow{g|_U} & \mathbb{P}^1 \setminus \{0, 1/256, \infty\}
 \end{array}$$

and $U = g^{-1}(\mathbb{P}^1 \setminus \{0, 1/256, \infty\})$.

Moreover, the preimage of 0 under g consists of either 1 or 2 points. If $g^{-1}(0)$ is 2 points, then g is ramified to order 1, 2 or 4 at each point in $g^{-1}(0)$. If $g^{-1}(0)$ is a single point, then g is ramified to order 8 at that point. If g is unramified over $1/256$ then W is smooth, but if g ramifies over $1/256$ then W may have isolated singularities.

Using this, in [DHNT16, Section 4] we obtained a classification of smooth deformation equivalent families of such Calabi-Yau threefolds, under the assumption that g is unramified over $1/256$ (which, by the theorem above, ensures that our Calabi-Yau threefolds will be smooth). This classification is determined by two pieces of data: a pair of numbers i and j in $\{1, 2, 4\}$, denoting the orders of ramification of g over 0, and a choice of partition $\mu := [x_1, \dots, x_k]$ of $\deg(g) = i + j$, denoting the ramification profile of g over ∞ . Note here that the case where $g^{-1}(0)$ is a single point is a deformation of the case where $i = j = 4$, so may be ignored; see [DHNT16, Remark 3.1]. We call a general member of this family $W_{i,j}^\mu$.

THEOREM 5.5. [DHNT16, Propositions 3.5, 3.8] *If $W_{i,j}^\mu$ is as above, then*

- (1) $h^{2,1}(W_{i,j}^\mu) = k$,
- (2) $h^{1,1}(W_{i,j}^\mu) = 20 + \sum_{s=1}^k (2x_s^2 + 1) + c_i + c_j$.

where $c_1 = 30$, $c_2 = 10$ and $c_4 = 0$.

Finally, the proof of [DHNT16, Proposition 2.5] classifies the singular fibres of these threefolds, up to small birational transformations. We give names to each possibility and a description of one member of each equivalence class.

- (1) I_0 : A smooth K3 surface. Along with the generic fibre, which clearly has type I_0 , if $t \in g^{-1}(0)$ and g ramifies to order 4 at t , then the fibre over t is of type I_0 .
- (2) I_{odp} : A K3 surface with a single node. If $t \in g^{-1}(1/256)$, then the fibre over t is of type I_{odp} .
- (3) I_n^Δ for $n \in \mathbb{N}$: this is a semistable singular fibre whose dual graph is a triangulation of the faces of a 3-dimensional simplex with sides of length n . Such a fibre has $2n^2 + 2$ irreducible components. If $t \in g^{-1}(\infty)$ and g ramifies to order n at t , then the fibre over t is of type I_n^Δ .
- (4) II: A union of 11 smooth rational surfaces, one of which, E , has multiplicity 2, and the others, F_1, \dots, F_{10} , have multiplicity 1. Each F_i intersects E in a smooth rational curve, but $F_i \cap F_j$ is empty for $i \neq j$. If $t \in g^{-1}(0)$ and g ramifies to order 2 at t , then the fibre over t is of type II.
- (5) IV: A normal crossings union of 31 smooth rational surfaces. One has multiplicity 4, and the rest have multiplicities 3, 2 or 1. If $t \in g^{-1}(0)$ and g is unramified at t , then the fibre over t is of type IV.

5.3. Degenerations in the mirror. Now we look at the mirror picture. The Dolgachev-Nikulin mirror to an M_2 -polarized K3 surface is a K3 surface of degree 4 ((4) -polarized); generically such K3 surfaces are just smooth quartics in \mathbb{P}^3 . There are three Fano threefolds which contain a K3 surface of degree 4 as their anticanonical hypersurface; these are X_1 , the quartic threefold in \mathbb{P}^4 , X_2 , the double cover of \mathbb{P}^3 ramified along a smooth generic quartic surface, and X_4 , which is just \mathbb{P}^3 itself. Note that the integer i assigned to each Fano X_i is just the index of that Fano threefold.

Let Z be a generic smooth anticanonical K3 surface in both X_i and X_j , for some choice of integers i, j in $\{1, 2, 4\}$. Then $N_{Z/X_i} \cong \omega_{X_i}^{-1}|_Z \cong \mathcal{O}_Z(i)$, where $\mathcal{O}_Z(1)$ is the restriction of a hyperplane section in \mathbb{P}^3 to Z . Thus if we take a normal crossings union of X_i and X_j meeting along Z then, in accordance with the discussion in Section 2.1, we cannot construct a Calabi-Yau threefold by smoothing $X_i \cup_Z X_j$, since $N_{Z/X_i} \otimes N_{Z/X_j} = \mathcal{O}_Z(i + j) \neq \mathcal{O}_Z$.

This is similar to the situation in Section 3.1, where we could not smooth $X_1 \cup_Z X_2$ without first blowing up the locus $V \cap X_1 \cap X_2$, and we will solve it in the same way. Let C_1, \dots, C_k be smooth curves in S cut out by sections of $\mathcal{O}_Z(x_s)$, for positive integers x_1, \dots, x_k , so that $\sum_{s=1}^k x_s = i + j$. Then let $f: \tilde{X}_i \rightarrow X_i$ be the blow up of X_i in C_1, \dots, C_k sequentially and let E_i be the exceptional divisor over C_i , for $i = 1, \dots, k$.

The canonical divisor of \tilde{X}_i is given by $f^*K_{X_i} - \sum_{s=1}^k E_s$, so $\omega_{\tilde{X}_i}^{-1} \cong \mathcal{O}_Z(-j)$. Therefore, according to [KN94, Theorem 4.2], we may smooth $\tilde{X}_i \cup_Z X_j$ to a Calabi-Yau threefold. We denote this threefold by $V_{i,j}^\mu$, where, as before, μ denotes the partition $[x_1, \dots, x_k]$ of $(i + j)$.

We claim that $V_{i,j}^\mu$ and $W_{i,j}^\mu$ are mirror dual, in the classical sense. As a first piece of evidence for this, we compute the Hodge numbers of $V_{i,j}^\mu$.

PROPOSITION 5.6. *Let $i, j \in \{1, 2, 4\}$ and let $\mu = [x_1, \dots, x_k]$ be a partition of $i + j$. Then the Hodge numbers of the threefold $V_{i,j}^\mu$ are given by*

$$h^{1,1}(V_{i,j}^\mu) = k,$$

$$h^{2,1}(V_{i,j}^\mu) = 20 + \sum_{s=1}^k (2x_s^2 + 1) + h^{2,1}(X_i) + h^{2,1}(X_j),$$

where $h^{2,1}(X_s) = 30$ (resp. 10, 0) for $s = 1$ (resp. 2, 4).

PROOF. By definition, $V_{i,j}^\mu$ is a smoothing of $\tilde{X}_i \cup_Z X_j$. Define

$$q := \text{rank}(\text{im}(H^2(\tilde{X}_i, \mathbb{Z}) \oplus H^2(X_j, \mathbb{Z}) \rightarrow H^2(Z, \mathbb{Z}))).$$

Then Lee [Lee10, Corollary 8.2] shows that the Hodge numbers of $V_{i,j}^\mu$ are given by

$$h^{1,1}(V_{i,j}^\mu) = h^2(\tilde{X}_i) + h^2(X_j) - q - 1,$$

$$h^{2,1}(V_{i,j}^\mu) = 21 + h^{2,1}(\tilde{X}_i) + h^{2,1}(X_j) - q.$$

Now, since the Néron-Severi group of Z is generated by the restriction of a hyperplane section from X_i , we must have $q = 1$. Moreover, since we blew up X_i a total of k times to obtain \tilde{X}_i , we have $h^2(\tilde{X}_i) = k + 1$ and $h^2(X_j) = 1$. Thus $h^{1,1}(V_{i,j}^\mu) = k$.

To compute $h^{2,1}(V_{i,j}^\mu)$, we begin by noting that a smooth curve C_s defined by a section of $\mathcal{O}_Z(x_s)$ has self-intersection $4x_s^2$ in Z . So the genus formula for curves on a surface gives $g(C_s) = 2x_s^2 + 1$. Thus, by standard results on the cohomology of a blow-up (see, for instance, [Voi07, Theorem 7.31]), we find

$$h^{2,1}(\tilde{X}_i) = h^{2,1}(X_i) + \sum_{s=1}^k (2x_s^2 + 1),$$

giving the claimed result for $h^{2,1}(V_{i,j}^\mu)$. Finally, the values of $h^{2,1}(X_s)$ are easy to compute explicitly. □

Putting this proposition together with the result of Theorem 5.5, we obtain:

COROLLARY 5.7. *Let $i, j \in \{1, 2, 4\}$ be a pair of integers and let $\mu = [x_1, \dots, x_k]$ be a partition of $i + j$. Then there is a mirror duality between the Hodge numbers of the Calabi-Yau threefolds $V_{i,j}^\mu$ and $W_{i,j}^\mu$.*

We expect that $V_{i,j}^\mu$ is actually mirror to $W_{i,j}^\mu$, but of course this is not a proof. As further evidence, however, we can also compare filtrations as in Section 5.1. For the threefolds $V_{i,j}^\mu$, we may compute the limit mixed Hodge structure associated to the degeneration to $\tilde{X}_i \cup_Z X_j$, to obtain

	Gr_F^3	Gr_F^2	Gr_F^1	Gr_F^0
Gr_4^M	\mathbb{C}	\mathbb{C}^{19}	\mathbb{C}	0
Gr_3^M	0	\mathbb{C}^v	\mathbb{C}^v	0
Gr_2^M	0	\mathbb{C}	\mathbb{C}^{19}	\mathbb{C}

for $v = h^{2,1}(X_i) + h^{2,1}(X_j) + \sum_{s=1}^k (2x_s^2 + 1)$.

Now, for the threefolds $W_{i,j}^\mu$, let J be the cup product operator with the negative of the class of a fibre of the M_2 -polarized K3 surface fibration on $W_{i,j}^\mu$. Then, by the proof of [DHNT16, Proposition 3.5] and Proposition 5.6, we see that the rank of the space of divisors in $H^{1,1}(W_{i,j}^\mu)$ that are supported on fibres is

$$\text{rank}(\ker(J)) = 1 + \sum_{s=1}^k (2x_s^2 + 1) + h^{2,1}(X_i) + h^{2,1}(X_j) = v + 1.$$

Moreover, the image of J in $H^{1,1}(W_{i,j}^\mu)$ is the span of the class of a fibre and the image of J in $H^{3,3}(W_{i,j}^\mu)$ spans $H^{3,3}(W_{i,j}^\mu)$. Finally, the image of J in $H^{2,2}(W_{i,j}^\mu)$ is the space of classes dual to divisors in $H^{1,1}(W_{i,j}^\mu)$ swept out by monodromy invariant divisors on a general fibre, which has rank 19. Thus, we obtain

	$H^{0,0}(X_{i,j}^\mu)$	$H^{1,1}(W_{i,j}^\mu)$	$H^{2,2}(W_{i,j}^\mu)$	$H^{3,3}(W_{i,j}^\mu)$
coimage(J)	\mathbb{C}	\mathbb{C}^{19}	\mathbb{C}	0
$\ker(J)/\text{im}(J)$	0	\mathbb{C}^v	\mathbb{C}^v	0
im(J)	0	\mathbb{C}	\mathbb{C}^{19}	\mathbb{C}

and the duality of bifiltered vector spaces discussed in Section 5.1 is satisfied in this case.

Finally, we note that restriction of divisors from X_i and X_j induces a lattice polarization of Z by the lattice $\langle 4 \rangle$, whilst the K3 surface fibration on $W_{i,j}^\mu$ is M_2 -polarized. As expected from our previous calculations, these two lattices are Dolgachev-Nikulin mirror dual.

5.4. Relationship to LG models. Now we will see how this fits with the results of Section 2. We begin by describing the LG models of X_1, X_2 and X_4 . Again, we note that these threefolds can be constructed from the basic K3 fibred threefold \mathcal{X} .

THEOREM 5.8. [Gol07, DHK16] *Define $g_\ell: \mathbb{A}^1 \rightarrow \mathbb{A}^1$ by $g_\ell(t) = t^\ell$. Then the LG model $(Y_\ell, \mathbf{w}_\ell)$ of X_ℓ is a partial compactification of $g_\ell^* \mathcal{X}$.*

In fact, it follows from our classification of singular fibres in Section 5.2 that

- (1) (Y_1, \mathbf{w}_1) has two singular fibres in \mathbb{A}^1 , located at 0 and $1/256$, of types IV and I_{odp} respectively,
- (2) (Y_2, \mathbf{w}_2) has three singular fibres in \mathbb{A}^1 , located at 0 and $\pm\sqrt{1/256}$, of types II and I_{odp} respectively,
- (3) (Y_4, \mathbf{w}_4) has four singular fibres in \mathbb{A}^1 , located at $\sqrt[4]{1/256}$, all of which have type I_{odp} . Its fibre over 0 has type I_0 .

The following ansatz has been suggested by Katzarkov [Kat09]:

ANSATZ 5.9. *Blowing up a Fano threefold X in a smooth curve of genus $g \geq 2$ has the effect of deforming the LG model (Y, \mathbf{w}) of X , so that a semistable fibre with $g + 1$ irreducible components moves from infinity to a point in \mathbb{A}^1 .*

By this logic, we can induce that if \tilde{X}_ℓ is as above, then its LG model $(\tilde{Y}_\ell, \tilde{\mathbf{w}}_\ell)$ has singular fibres of the types listed above for $(Y_\ell, \mathbf{w}_\ell)$, along with fibres of type $I_{x_s}^\Delta$, for $s = 1, \dots, k$, since a smooth curve C_s determined by a section of $\mathcal{O}_Z(x_s)$ has genus $2x_s^2 + 1$.

By the results of [DHNT16, Section 4], generically we may choose the map g determining $W_{i,j}^\mu$, so that its ramification points not lying over 0 and ∞ are all of order 2, and so that all nonzero branch points λ of g have $|\lambda| \geq R$ for some real number $R > 1/256$. Then there are two components to the preimage of the disc $U_R = \{z \in \mathbb{C} : |z| < R\}$, corresponding to the two preimages of 0. Let U_R^ℓ be the component of $g^{-1}(U_R)$ containing the ramification point of g of order ℓ over 0 (for $\ell \in \{i, j\}$). Then the restriction of the fibration π on $W_{i,j}^\mu$ to U_R^ℓ is a fibration over a disc with a fibre of type IV, II or I_0 over $g^{-1}(0)$, depending on whether $\ell = 1, 2$ or 4 respectively, and ℓ fibres of type I_{odp} . In fact, this fibration over U_R^ℓ is deformation equivalent to the LG model of X_ℓ , in the sense that as $R \rightarrow \infty$, the map g degenerates to a stable map from a pair of rational curves C_i and C_j meeting at a point, such that the restriction of g to C_ℓ is g_ℓ . We also see that monodromy around the boundary of U_R^ℓ is equal to the monodromy around ∞ of the the LG model $(Y_\ell, \mathbf{w}_\ell)$.

Away from 0 and ∞ , we have that the restriction of π to $g^{-1}(\mathbb{P}^1 \setminus U_R)$ has singular fibres of type $I_{x_i}^\Delta$, for $i = 1, \dots, k$, and these account for all of the singular fibres of π restricted to $g^{-1}(\mathbb{P}^1 \setminus U_R)$. We thus obtain the following theorem; which is highly reminiscent of our philosophy in the K3 surface case (see Section 4.3).

THEOREM 5.10. *The threefold $W_{i,j}^\mu$ is topologically equivalent to the gluing of the LG model of X_j to the LG model of \tilde{X}_i , as described in Section 2.2.*

6. Non-commutative fibrations

Now we turn our attention to ways in which this construction can fail. Suppose that V and W are mirror dual Calabi-Yau threefolds. The computation in

Section 5.1 suggests that, if a Tyurin degeneration of V occurs along a locus in the moduli space of V that contains a maximally unipotent monodromy point, then there should exist a K3 fibration on W whose properties are consistent with the results in Section 2.3.

However, it is also possible for Tyurin degenerations to occur along loci in the moduli space of V that are disjoint from points of maximally unipotent monodromy. For such Tyurin degenerations the argument in Section 5.1 will break down and, in particular, in such cases we will have no guarantee of the existence of K3 fibrations on W . For an example where this occurs, consider:

EXAMPLE 6.1. Let W be the complete intersection of two cubics in \mathbb{P}^5 and let V be its Batyrev-Borisov mirror. Since $h^{2,1}(V) = h^{1,1}(W) = 1$, the dimension of the moduli space of V is 1 and, indeed, it can be shown that the moduli space of Calabi-Yau varieties deformation equivalent to V is $\mathbb{P}^1 \setminus \{0, 1, \infty\}$. In [KP09] Katzarkov and Przyjalkowski show that, after semi-stabilization, the degeneration of V over ∞ is a Tyurin degeneration. The degenerate fibre is a union of two quasi-Fano threefolds X_1, X_2 meeting in a K3 surface Z , which has Picard rank 20 and transcendental lattice isomorphic to

$$\begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}.$$

There are several problems with this example. First of all, we find that this Tyurin degeneration is not connected to a point of maximally unipotent monodromy in the moduli space of V , so the discussion above does not hold. An even more grave issue is the fact that the K3 surface Z has no Dolgachev-Nikulin mirror. We would therefore not expect a K3 surface fibration corresponding to this Tyurin degeneration to appear on the mirror Calabi-Yau threefold W and, indeed, the non-existence of such a fibration is easily verified, since $h^{1,1}(W) = 1$.

However, if we are willing to replace the mirror Calabi-Yau threefold W with its bounded derived category of coherent sheaves $\mathbf{D}^b(W)$, we *do* know how to apply mirror symmetry to monodromy in non-maximally unipotent families. Indeed, if \mathcal{V} is a family of Calabi-Yau threefolds over a punctured disc U , such that there is a symplectic form ω on \mathcal{V} which restricts to a symplectic form on each fibre, then the action of monodromy around 0 induces a symplectomorphism on a smooth fibre V . According to [Sze04], this symplectomorphism induces an autoequivalence on the Fukaya category $\mathcal{Fuk}(V, \omega)$, which passes through mirror symmetry to produce an element of $\text{AutEq}(\mathbf{D}^b(W))$.

Our goal is to understand what sort of autoequivalence this is. If $\pi: W \rightarrow \mathbb{P}^1$ is a fibration of W by K3 surfaces, then there is a right derived pullback functor $\mathbf{R}\pi^*: \mathbf{D}^b(\mathbb{P}^1) \rightarrow \mathbf{D}^b(W)$. We conjecture that the autoequivalence of $\mathbf{D}^b(W)$ obtained above is an autoequivalence related to the spherical functor $\mathbf{R}\pi^*$. Moreover, if p is a point in \mathbb{P}^1 , then the inclusion gives a pullback $\mathbf{R}i^*: \mathbf{D}^b(\mathbb{P}^1) \rightarrow \mathbf{D}^b(p)$. We can take the fibre product of these categories, denoted $\mathbf{D}^b(W) \otimes_{\mathbf{D}^b(\mathbb{P}^1)} \mathbf{D}^b(p)$, to give a category associated to a general fibre of W . We want this category to be the bounded derived category of coherent sheaves on a K3 surface.

To state this idea more precisely, we need a definition. A triangulated category \mathbf{T} with Serre functor \mathbf{S} is called *d-Calabi-Yau* if \mathbf{S} is equivalent to $[d]$, where $[d]$ indicates the natural “shift by d ” functor. This is a useful notion because the

bounded derived category $\mathbf{D}^b(X)$ of coherent sheaves on a smooth projective variety X is d -Calabi-Yau if and only if X is a Calabi-Yau variety of dimension d .

CONJECTURE 6.2. *Let V be a Calabi-Yau variety of dimension d that admits a Tyurin degeneration, and let W be a homological mirror of V . Then there is a functor $F: \mathbf{D}^b(\mathbb{P}^1) \rightarrow \mathbf{D}^b(W)$ so that for a generic point $p \in \mathbb{P}^1$, the category $\mathbf{D}^b(V) \otimes_{\mathbf{D}^b(\mathbb{P}^1)} \mathbf{D}^b(p)$ is a $(d - 1)$ -Calabi-Yau category.*

Moreover, we expect that the fibres of such a categorical fibration are equivalent, in an appropriate sense, to the Fukaya category of the Calabi-Yau $Z = X_1 \cap X_2$, where $X_1 \cup_Z X_2$ is the Tyurin degeneration of V .

EXAMPLE 6.3. Calabrese and Thomas [CT16] showed how this should work in the setting of Example 6.1. Let f_1 and f_2 be cubics so that $W = \{f_1 = f_2 = 0\} \subset \mathbb{P}^5$. Blowing up \mathbb{P}^5 along W , one obtains a five-fold $\tilde{\mathbb{P}}^5$ which is fibred over \mathbb{P}^1 by cubics $sf_1 + tf_2 = 0$, for $[s : t] \in \mathbb{P}^1$. This variety has derived category

$$\mathbf{D}^b(\tilde{\mathbb{P}}^5) = \langle \mathbf{D}^b(W), \tilde{\pi}^* \mathbf{D}^b(\mathbb{P}^1)(i, 0) : i = 3, 4, 5 \rangle$$

where $\tilde{\pi}: \tilde{\mathbb{P}}^5 \rightarrow \mathbb{P}^1$ is the natural map.

Using this, Calabrese and Thomas show that there is a fibration structure on $\mathbf{D}^b(W)$ satisfying the conditions of Conjecture 6.2 above. In this case the derived categories $\mathbf{D}^b(W) \otimes_{\mathbf{D}^b(\mathbb{P}^1)} \mathbf{D}^b(p)$ are 2-Calabi-Yau categories, but they are not the bounded derived category of coherent sheaves on any K3 or abelian surface; instead they are semi-orthogonal summands of the derived categories of the cubic fourfolds cut out by the equations $sf_1 + tf_2 = 0$.

Moreover, according to Hassett [Has99], the transcendental lattice of a generic cubic fourfold is isomorphic to

$$E_8^2 \oplus U^2 \oplus \begin{pmatrix} -2 & -1 \\ -1 & -2 \end{pmatrix},$$

which is precisely the Dolgachev-Nikulin dual lattice associated to the transcendental lattice of the K3 surface Z in Example 6.1. We thus conjecture that the non-commutative fibration found by Calabrese and Thomas [CT16] is mirror dual to the Tyurin degeneration of Katzarkov and Przyjalkowski [KP09].

It may be possible to generate other examples like this one as follows. Suppose that \mathcal{V} is a family of Calabi-Yau threefolds over a small disc U , so that the general fibre V has $h^{2,1}(V) = 1$ and the fibre over 0 is a Tyurin degeneration. Then the matrix describing the monodromy action on $H^3(V, \mathbb{Q})$ associated to a loop around 0 has two Jordan blocks of rank 1.

The families of Calabi-Yau variations of Hodge structure lying over $\mathbb{P}^1 \setminus \{0, 1, \infty\}$ with appropriate monodromy properties have been classified by Doran and Morgan [DM06]. One can check that there are precisely three families in this classification for which the monodromy matrix associated to ∞ has two Jordan blocks of rank 1. These families are mirrors to

- (1) The (3, 3) complete intersection in \mathbb{P}^5 ,
- (2) The (4, 4) complete intersection in $\mathbb{W}\mathbb{P}(1, 1, 1, 1, 2, 2)$,
- (3) The (6, 6) complete intersection in $\mathbb{W}\mathbb{P}(1, 1, 2, 2, 3, 3)$.

This raises the following natural question.

QUESTION 6.4. Are there categorical fibrations on $\mathbf{D}^b(W)$ satisfying Conjecture 6.2, for W a $(4, 4)$ complete intersection in $\mathbb{WP}(1, 1, 1, 1, 2, 2)$ or a $(6, 6)$ complete intersection in $\mathbb{WP}(1, 1, 2, 2, 3, 3)$?

Neither of these Calabi-Yau threefolds W can admit commutative K3 fibrations, since $h^{1,1}(W) = 1$, so if mirror dual fibrations exist then they are necessarily non-commutative, as in Example 6.3 above.

References

- [ACG13] Dan Abramovich, Qile Chen, Danny Gillam, Yuhao Huang, Martin Olsson, Matthew Satriano, and Shenghao Sun, *Logarithmic geometry and moduli*, Handbook of moduli. Vol. I, Adv. Lect. Math. (ALM), vol. 24, Int. Press, Somerville, MA, 2013, pp. 1–61. MR3184161
- [AKMS97] A. C. Avram, M. Kreuzer, M. Mandelberg, and H. Skarke, *Searching for K3 fibrations*, Nuclear Phys. B **494** (1997), no. 3, 567–589, DOI 10.1016/S0550-3213(97)00214-9. MR1453773
- [Ale96a] V. Alexeev, *Log canonical singularities and complete moduli of stable pairs*, Preprint, August 1996, [arXiv:alg-geom/9608013](https://arxiv.org/abs/alg-geom/9608013).
- [Ale96b] ———, *Moduli spaces $M_{g,n}(W)$ for surfaces*, Higher-Dimensional Complex Varieties (Trento, 1994) (Berlin), de Gruyter, 1996, pp. 1–22.
- [Ale06] ———, *Higher-dimensional analogues of stable curves*, International Congress of Mathematicians. Vol. II, Eur. Math. Soc., Zürich, 2006, pp. 515–536.
- [AT16] V. Alexeev and A. Thompson, *Modular compactification of moduli of K3 surfaces of degree 2*, Manuscript in preparation, 2016, preliminary version available at <http://alpha.math.uga.edu/~valery/>.
- [Aur08] Denis Auroux, *Special Lagrangian fibrations, mirror symmetry and Calabi-Yau double covers* (English, with English and French summaries), Astérisque **321** (2008), 99–128. Géométrie différentielle, physique mathématique, mathématiques et société. I. MR2521645
- [Bat94] Victor V. Batyrev, *Dual polyhedra and mirror symmetry for Calabi-Yau hypersurfaces in toric varieties*, J. Algebraic Geom. **3** (1994), no. 3, 493–535. MR1269718
- [BB96] Victor V. Batyrev and Lev A. Borisov, *On Calabi-Yau complete intersections in toric varieties*, Higher-dimensional complex varieties (Trento, 1994), de Gruyter, Berlin, 1996, pp. 39–65. MR1463173
- [Bor93] L. A. Borisov, *Towards the mirror symmetry for Calabi-Yau complete intersections in Gorenstein toric Fano varieties*, Preprint, October 1993, [arXiv:alg-geom/9310001](https://arxiv.org/abs/alg-geom/9310001).
- [Bru15] A. Brunyate, *A compact moduli space of elliptic K3 surfaces*, Ph.D. thesis, University of Georgia, 2015.
- [CCS13] Philip Candelas, Andrei Constantin, and Harald Skarke, *An abundance of K3 fibrations from polyhedra with interchangeable parts*, Comm. Math. Phys. **324** (2013), no. 3, 937–959, DOI 10.1007/s00220-013-1802-2. MR3123541
- [CD07] Adrian Clingher and Charles F. Doran, *Modular invariants for lattice polarized K3 surfaces*, Michigan Math. J. **55** (2007), no. 2, 355–393, DOI 10.1307/mmj/1187646999. MR2369941
- [CK99] David A. Cox and Sheldon Katz, *Mirror symmetry and algebraic geometry*, Mathematical Surveys and Monographs, vol. 68, American Mathematical Society, Providence, RI, 1999. MR1677117
- [CLS11] David A. Cox, John B. Little, and Henry K. Schenck, *Toric varieties*, Graduate Studies in Mathematics, vol. 124, American Mathematical Society, Providence, RI, 2011. MR2810322
- [CT16] John R. Calabrese and Richard P. Thomas, *Derived equivalent Calabi-Yau threefolds from cubic fourfolds*, Math. Ann. **365** (2016), no. 1-2, 155–172, DOI 10.1007/s00208-015-1260-6. MR3498907
- [DHK16] C. F. Doran, A. Harder, L. Katzarkov, J. Lewis, and V. Przyjalkowski, *Moduli spaces of K3 surfaces and mirror symmetry for Fano threefolds*, Manuscript in preparation, 2016.

- [DHNT15] Charles F. Doran, Andrew Harder, Andrey Y. Novoseltsev, and Alan Thompson, *Families of lattice polarized K3 surfaces with monodromy*, Int. Math. Res. Not. IMRN **23** (2015), 12265–12318, DOI 10.1093/imrn/rnv071. MR3431621
- [DHNT16] C. F. Doran, A. Harder, A. Y. Novoseltsev, and A. Thompson, *Calabi-Yau threefolds fibred by mirror quartic K3 surfaces*, Adv. Math. **298** (2016), 369–392, DOI 10.1016/j.aim.2016.03.045. MR3505744
- [DM06] Charles F. Doran and John W. Morgan, *Mirror symmetry and integral variations of Hodge structure underlying one-parameter families of Calabi-Yau threefolds*, Mirror symmetry. V, AMS/IP Stud. Adv. Math., vol. 38, Amer. Math. Soc., Providence, RI, 2006, pp. 517–537. MR2282973
- [Dol96] I. V. Dolgachev, *Mirror symmetry for lattice polarized K3 surfaces*, J. Math. Sci. **81** (1996), no. 3, 2599–2630, DOI 10.1007/BF02362332. Algebraic geometry, 4. MR1420220
- [EHX97] Tohru Eguchi, Kentaro Hori, and Chuan-Sheng Xiong, *Gravitational quantum cohomology*, Internat. J. Modern Phys. A **12** (1997), no. 9, 1743–1782, DOI 10.1142/S0217751X97001146. MR1439892
- [Fri83] Robert Friedman, *Global smoothings of varieties with normal crossings*, Ann. of Math. (2) **118** (1983), no. 1, 75–114, DOI 10.2307/2006955. MR707162
- [Giv98] Alexander Givental, *A mirror theorem for toric complete intersections*, Topological field theory, primitive forms and related topics (Kyoto, 1996), Progr. Math., vol. 160, Birkhäuser Boston, Boston, MA, 1998, pp. 141–175. MR1653024
- [Gol07] Vasily V. Golyshev, *Classification problems and mirror duality*, Surveys in geometry and number theory: reports on contemporary Russian mathematics, London Math. Soc. Lecture Note Ser., vol. 338, Cambridge Univ. Press, Cambridge, 2007, pp. 88–121, DOI 10.1017/CBO9780511721472.004. MR2306141
- [GP13] Antonella Grassi and Vittorio Perduca, *Weierstrass models of elliptic toric K3 hypersurfaces and symplectic cuts*, Adv. Theor. Math. Phys. **17** (2013), no. 4, 741–770. MR3262515
- [GS75] Phillip Griffiths and Wilfried Schmid, *Recent developments in Hodge theory: a discussion of techniques and results*, Discrete subgroups of Lie groups and applicatons to moduli (Internat. Colloq., Bombay, 1973), Oxford Univ. Press, Bombay, 1975, pp. 31–127. MR0419850
- [Har16] A. Harder, *The geometry of Landau-Ginzburg models*, Ph.D. thesis, University of Alberta, 2016.
- [Has99] Brendan Hassett, *Some rational cubic fourfolds*, J. Algebraic Geom. **8** (1999), no. 1, 103–114. MR1658216
- [Kat92] Sheldon Katz, *Rational curves on Calabi-Yau threefolds*, Essays on mirror manifolds, Int. Press, Hong Kong, 1992, pp. 168–180. MR1191424
- [Kat09] L. Katzarkov, *Homological mirror symmetry and algebraic cycles*, Homological mirror symmetry, Lecture Notes in Phys., vol. 757, Springer, Berlin, 2009, pp. 125–152. MR2596637
- [KKP14] Ludmil Katzarkov, Maxim Kontsevich, and Tony Pantev, *Bogomolov-Tian-Todorov theorems for Landau-Ginzburg models*, J. Differential Geom. **105** (2017), no. 1, 55–117. MR3592695
- [KN94] Yujiro Kawamata and Yoshinori Namikawa, *Logarithmic deformations of normal crossing varieties and smoothing of degenerate Calabi-Yau varieties*, Invent. Math. **118** (1994), no. 3, 395–409, DOI 10.1007/BF01231538. MR1296351
- [KP09] L. Katzarkov and V. Przyjalkowski, *Generalized homological mirror symmetry and cubics*, Tr. Mat. Inst. Steklova **264** (2009), no. Mnogomernaya Algebraicheskaya Geometriya, 94–102, DOI 10.1134/S0081543809010118; English transl., Proc. Steklov Inst. Math. **264** (2009), no. 1, 87–95. MR2590839
- [KSB88] J. Kollár and N. I. Shepherd-Barron, *Threefolds and deformations of surface singularities*, Invent. Math. **91** (1988), no. 2, 299–338, DOI 10.1007/BF01389370. MR922803
- [Lee06] Nam-Hoon Lee, *Constructive Calabi-Yau manifolds*, ProQuest LLC, Ann Arbor, MI, 2006. Thesis (Ph.D.)—University of Michigan. MR2708981
- [Lee10] Nam-Hoon Lee, *Calabi-Yau construction by smoothing normal crossing varieties*, Internat. J. Math. **21** (2010), no. 6, 701–725, DOI 10.1142/S0129167X10006173. MR2658406

- [Mor84] David R. Morrison, *The Clemens-Schmid exact sequence and applications*, Topics in transcendental algebraic geometry (Princeton, N.J., 1981/1982), Ann. of Math. Stud., vol. 106, Princeton Univ. Press, Princeton, NJ, 1984, pp. 101–119. MR756848
- [Ogu01] Keiji Oguiso, *On the finiteness of fiber-space structures on a Calabi-Yau 3-fold*, J. Math. Sci. (New York) **106** (2001), no. 5, 3320–3335, DOI 10.1023/A:1017959510524. Algebraic geometry, 11. MR1878052
- [Orl09] Dmitri Orlov, *Derived categories of coherent sheaves and triangulated categories of singularities*, Algebra, arithmetic, and geometry: in honor of Yu. I. Manin. Vol. II, Progr. Math., vol. 270, Birkhäuser Boston, Inc., Boston, MA, 2009, pp. 503–531, DOI 10.1007/978-0-8176-4747-6_16. MR2641200
- [PS08] Chris A. M. Peters and Joseph H. M. Steenbrink, *Mixed Hodge structures*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], vol. 52, Springer-Verlag, Berlin, 2008. MR2393625
- [Roh04] F. Rohsiepe, *Lattice polarized toric K3 surfaces*, Preprint, September 2004, [arXiv:hep-th/0409290](https://arxiv.org/abs/hep-th/0409290).
- [Sca87] Francesco Scattone, *On the compactification of moduli spaces for algebraic K3 surfaces*, Mem. Amer. Math. Soc. **70** (1987), no. 374, x+86, DOI 10.1090/memo/0374. MR912636
- [Sei01a] Paul Seidel, *More about vanishing cycles and mutation*, Symplectic geometry and mirror symmetry (Seoul, 2000), World Sci. Publ., River Edge, NJ, 2001, pp. 429–465, DOI 10.1142/9789812799821_0012. MR1882336
- [Sei01b] Paul Seidel, *Vanishing cycles and mutation*, European Congress of Mathematics, Vol. II (Barcelona, 2000), Progr. Math., vol. 202, Birkhäuser, Basel, 2001, pp. 65–85. MR1905352
- [Sze04] Balázs Szendrői, *Contractions and monodromy in homological mirror symmetry*, Strings and geometry, Clay Math. Proc., vol. 3, Amer. Math. Soc., Providence, RI, 2004, pp. 301–314. MR2103728
- [Tyu04] Andrei N. Tyurin, *Fano versus Calabi-Yau*, The Fano Conference, Univ. Torino, Turin, 2004, pp. 701–734. MR2112600
- [Voi07] Claire Voisin, *Hodge theory and complex algebraic geometry. II*, Reprint of the 2003 English edition, Cambridge Studies in Advanced Mathematics, vol. 77, Cambridge University Press, Cambridge, 2007. Translated from the French by Leila Schneps. MR2449178

DEPARTMENT OF MATHEMATICAL AND STATISTICAL SCIENCES, 632 CAB, UNIVERSITY OF ALBERTA, EDMONTON, AB, T6G 2G1, CANADA

E-mail address: charles.doran@ualberta.ca

DEPARTMENT OF MATHEMATICAL AND STATISTICAL SCIENCES, 632 CAB, UNIVERSITY OF ALBERTA, EDMONTON, AB, T6G 2G1, CANADA

E-mail address: aharder@ualberta.ca

DEPARTMENT OF PURE MATHEMATICS, UNIVERSITY OF WATERLOO, 200 UNIVERSITY AVE WEST, WATERLOO, ON, N2L 3G1, CANADA

E-mail address: am6thomp@uwaterloo.ca